

Regimes of stratified turbulence at low Prandtl number

Kasturi Shah^{1,2}, Gregory P. Chini^{3,4}, Colm-cille P. Caulfield^{1,5} and Pascale Garaud⁶

¹ Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge CB3 0WA, UK

² Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge MA 02139, USA

³ Program in Integrated Applied Mathematics, University of New Hampshire, Durham, NH 03824, USA.

⁴ Department of Mechanical Engineering, University of New Hampshire, Durham, NH 03824, USA

⁵ Institute for Energy and Environmental Flows, University of Cambridge, Cambridge CB3 0EZ, UK

⁶ Department of Applied Mathematics, Baskin School of Engineering, University of California Santa Cruz, Santa Cruz, CA 95064, USA

Abstract

Quantifying transport by strongly stratified turbulence in low Prandtl number (Pr) fluids is critically important for the development of better models for the structure and evolution of stellar interiors. Motivated by recent numerical simulations showing strongly anisotropic flows suggestive of scale-separated dynamics, we perform a multiscale asymptotic analysis of the governing equations. We find that, in all cases, the resulting slow-fast system naturally takes a quasilinear form. Our analysis also reveals the existence of several distinct dynamical regimes depending on the emergent buoyancy Reynolds and Péclet numbers, $Re_b = \alpha^2 Re$ and $Pe_b = Pr Re_b$, respectively, where α is the aspect ratio of the large-scale turbulent flow structures, and Re is the outer scale Reynolds number. Scaling relationships relating the aspect ratio, the characteristic vertical velocity, and the strength of the stratification (measured by the Froude number Fr) naturally emerge from the analysis. When $Pe_b \ll \alpha$, the dynamics at all scales is dominated by buoyancy diffusion, and our results recover the scaling laws empirically obtained from direct numerical simulations by Cope et al. (2020). For $Pe_b > O(1)$, diffusion is negligible (or at least subdominant) at all scales and our results are consistent with those of Chini et al. (2022) for strongly stratified geophysical turbulence at $Pr = O(1)$. Finally, we have identified a new regime for $\alpha \ll Pe_b \ll 1$, in which slow, large scales are diffusive while fast, small scales are not. We conclude by presenting a map of parameter space that clearly indicates the transitions between isotropic turbulence, non-diffusive stratified turbulence, diffusive stratified turbulence and viscously-dominated flows.

1 Introduction

Quantifying vertical transport (of heat, chemical tracers and momentum) in the stably stratified regions of stellar interiors is paramount to a better understanding of their structure and dynamics. Specification of this transport has long been a significant source of uncertainty in evolution models of stars, as the vertical transport rates required to match observations are largely inconsistent with purely diffusive processes. The same problem affects our attempts to model stably stratified regions of the Earth's atmosphere and oceans. The solution in both cases is to include turbulent transport in the models, but this in turn requires reliable parametrizations of small-scale fluxes as functions of the local properties of the fluid and of its large-scale flow (Munk, 1966; Pinsonneault, 1997; Ivey et al., 2008; Gregg et al., 2018; Aerts et al., 2019; Caulfield, 2021).

Stratified turbulence is intrinsically complex. Indeed, relatively small-scale turbulent vertical mixing in a stably stratified region typically incurs an energetic cost to lift dense fluid parcels up irreversibly and correspondingly lower buoyant parcels down; as such it can only be sustained on long time scales by tapping into some larger-scale energy reservoir. Different fluid instabilities access different sources of energy: for instance, baroclinic and double-diffusive instabilities draw from the available potential energy of the fluid, magnetic instabilities rely on the magnetic energy of a finite-amplitude large-scale field, and shear instabilities tap into the kinetic energy reservoir of a larger-scale mean flow. The nature of the turbulence driven by each

of these instabilities differs substantially, so that it “still remains extremely difficult to say anything generic about [stratified] mixing” (Caulfield, 2021).

In this work, we focus on modeling turbulence in stellar interiors driven exclusively by shear instabilities. Stellar fluids generally have a low Prandtl number Pr , where

$$Pr = \frac{\nu^*}{\kappa^*}, \quad (1)$$

ν^* is the kinematic viscosity of the fluid, and κ^* is the buoyancy diffusivity (e.g. thermal or compositional diffusivity, depending on the main source of stratification); see Figure 7 of Garaud et al. (2015b). Note that in all that follows, dimensional quantities are starred, while non-dimensional quantities are not. We ignore rotation, magnetic fields, and the presence of multiple components contributing to the flow buoyancy. Although these effects undoubtedly are important in most situations, the dynamics of purely shear-driven stratified turbulence in low Pr fluids is still far from well understood, justifying the narrow scope of this study.

Naively, one might expect a *vertical* shear to be the most natural source of *vertical* mixing in stably stratified fluids. Balancing the potential energy cost and the kinetic energy gain of adiabatic turbulent eddies in a vertically sheared flow, Richardson (1920) concluded that turbulence can be sustained provided

$$J = \frac{N^{*2}}{S^{*2}} \leq O(1), \quad (2)$$

where N^* is the typical value of the buoyancy frequency of the stratification, and S^* is the typical vertical shearing rate of the mean flow. Condition (2) is known today as the Richardson criterion (where the nondimensional parameter J is commonly referred to as a gradient Richardson number, which can vary with the vertical position z^*). For linear normal mode disturbances in an inviscid, non-diffusive parallel shear flow, Miles (1961) and Howard (1961) formalized this argument to establish rigorously the necessary condition for linear instability to be that $J(z^*) < 1/4$ somewhere within the flow. In the stably stratified regions of the ocean, there is increasing evidence that flows often are marginally stable, with the minimum value of $J(z^*) \simeq 1/4$ (Smyth and Moum, 2013), suggestive of ‘self-organized criticality’ (Salehipour et al., 2018; Smyth et al., 2019; Mashayek et al., 2022). This criterion seems to be rarely satisfied in stars, however, because J is usually very large (see, e.g. Garaud, 2021) except in regions where the stratification itself is exceptionally weak.

The Richardson criterion can be relaxed when the flow disturbances are not purely adiabatic, because the energy cost of vertical motions is reduced if the advected fluid parcels radiatively (Townsend, 1958; Dudas, 1973) or diffusively (Zahn, 1974; Jones, 1977; Lignières et al., 1999) adjust their buoyancy to that of the background stratification on a time scale comparable with, or shorter than, their turnover time scale. In the diffusive case, this adjustment is sufficiently fast when the eddy Péclet number

$$Pe_\ell \equiv \frac{u_\ell^* \ell^*}{\kappa^*} \leq O(1), \quad (3)$$

where u_ℓ^* is the characteristic velocity of an eddy of size ℓ^* . Zahn (1974) heuristically argued that an appropriate criterion for the instability of diffusive eddies of size ℓ^* in a vertical shear flow is

$$JPe_\ell \leq O(1), \quad (4)$$

provided (3) holds. According to his criterion, it ought to be possible to maintain turbulence in very strongly stratified shear flows ($J \gg 1$) provided eddies are small enough to ensure that $Pe_\ell \leq O(J^{-1})$.

Note that these so-called ‘diffusive’ vertical shear instabilities are irrelevant in the context of oceanic and atmospheric mixing, because the Prandtl number of air or water under geophysically relevant conditions is usually of order unity or much larger. As a result, the condition $Pe_\ell \leq O(1)$ equivalently implies that $Re_\ell \equiv u_\ell^* \ell^* / \nu^* \leq O(1)$, where Re_ℓ is the corresponding eddy Reynolds number. As shear instabilities must have a relatively large Reynolds number to develop (otherwise viscous energy losses are too great), the requirement that $Re_\ell \gg 1$ is incompatible with $Pe_\ell = Pr Re_\ell \leq O(1)$ when $Pr \geq O(1)$.

The situation is quite different in stellar interiors, as first noted by Spiegel and Zahn (1970) and Zahn (1974), because of their intrinsically small Prandtl number. With $Pr \ll 1$, and regardless of the size of the

outer scale Reynolds and Péclet numbers

$$Re \equiv \frac{U^*L^*}{\nu^*} \text{ and } Pe \equiv \frac{U^*L^*}{\kappa^*}, \quad (5)$$

(where U^* and L^* are the system-scale characteristic flow velocity and length scale, respectively), there is always an intermediate range of scales where turbulent eddies satisfy $Pe_\ell \ll 1 \ll Re_\ell$, namely, where diffusion dominates their dynamics while viscous forces remain negligible. This is a fundamental property of low Pr fluids that has important consequences for various other fluid instabilities as well (cf. the review by Garaud, 2021). In the case of vertical shear instabilities, this property naturally allows for the maintenance of diffusive stratified turbulence provided the shear is large enough for (4) to hold.

Even under this relaxed criterion, however, stellar interiors are so strongly stratified that their mean vertical shear is rarely unstable. Furthermore, in the few instances where it is, the resulting turbulent vertical transport is not much larger than the purely diffusive transport (Garaud, 2021). Therefore, it appears that a mean vertical shear cannot be a substantial and sustained source of turbulence in stellar interiors.

Meanwhile, a steadily growing body of evidence from laboratory experiments (Ruddick et al., 1989; Park et al., 1994; Holford and Linden, 1999; Oglethorpe et al., 2013; Jackson and Rehmann, 2014) and numerical simulations (Jacobitz and Sarkar, 2000; Basak and Sarkar, 2006; Brethouwer et al., 2007; Maffioli and Davidson, 2016; Lucas et al., 2017; Zhou and Diamessis, 2019; Yi and Koseff, 2023) is offering an alternative, much less-intuitive solution, namely that *horizontal* shear instabilities, or otherwise forced horizontal turbulence, can be a source of substantial and sustained *vertical* mixing in strongly stratified flows. In what follows, we review what is known about vertical mixing driven by horizontal flows, first at $Pr \geq O(1)$, where most of the research on this topic has been focused, and then at $Pr \ll 1$.

1.1 Stratified turbulence driven by horizontal flows at $Pr \geq O(1)$

Studies of stratified turbulence driven by horizontal flows generally quantify the effects of stratification using the outer-scale Froude number, defined as

$$Fr = \frac{U^*}{N^*L^*}. \quad (6)$$

In this expression U^* and L^* are now specifically assumed to be the characteristic *horizontal* velocity and length scale at the outer scale. Ruddick et al. (1989), Park et al. (1994) and Holford and Linden (1999) demonstrated experimentally that regular horizontal motions of vertical rods in a salt-stratified fluid (so $Pr = O(1000)$) with an initially constant stratification (characterized by a buoyancy frequency N^*) can cause substantial vertical mixing even when $Fr \ll 1$. Furthermore, they often observed the formation of steps in the density profile, with a characteristic vertical height $H^* \propto U^*/N^*$. Oglethorpe et al. (2013) similarly found substantial mixing and the formation of vertical layers in strongly stratified Taylor-Couette flow experiments, where the mean flow is by construction primarily horizontal.

Direct numerical simulations (DNS) are arguably more practical for investigation of $Pr = O(1)$ fluids (such as thermally-stratified air and water), and over the last 20 years significant progress in quantifying stratified turbulence in this parameter regime has been enabled by advances in supercomputing. In particular, access to the full three-dimensional structure of the velocity and buoyancy fields has provided new insights into the nature of turbulence at very large Reynolds numbers and very strong stratification, a regime characteristic of stratified turbulence in the ocean and atmosphere. Pioneering work by Brethouwer et al. (2007), for instance, has demonstrated that the eddy field is highly anisotropic in this regime, with a vertical eddy scale once again proportional to U^*/N^* – suggesting that this emergent ‘layer’ length scale is a universal property of stratified turbulence at both moderate and high Pr at sufficiently high Reynolds number, as theoretically predicted and experimentally demonstrated by Billant and Chomaz (2000); Billant and Chomaz (2001) (see also Caulfield, 2021, for a review).

Using the insight gained from their DNS data, Brethouwer et al. (2007) proposed an anisotropic rescaling of the governing equations, introducing the small vertical scale $H^* = FrL^* = U^*/N^*$ and taking the asymptotic limit $Fr \rightarrow 0$. Crucially, the horizontal scales are assumed to remain $O(L^*)$ in their work.

Inspection of the rescaled equations reveals the fundamental role of the so-called buoyancy Reynolds number

$$Re_b \equiv \left(\frac{H^*}{L^*} \right)^2 Re, \quad (7)$$

which needs to be substantially greater than one for viscous effects to be negligible (see Brethouwer et al., 2007; Bartello and Tobias, 2013, and § 2 for further details). In that case, the characteristic vertical velocities and buoyancy fluctuations are predicted to scale as $W^* \propto FrU^*$ and $B^* \propto FrL^*N^{*2} = H^*N^{*2}$, respectively.

Fundamental to the arguments presented in Brethouwer et al. (2007) is the concept that there is a single (large, outer) horizontal scale of significance, with the implicit consequence that spatio-temporal variability on smaller horizontal scales does not exert a controlling influence on the dynamics. This supposition is not entirely self-consistent, however, as Brethouwer et al. (2007) demonstrated that an inevitable consequence of their scaling is that $J \simeq O(1)$ at least somewhere in the flow, thus implying the possible existence of local shear instabilities that would drive fast, small-scale isotropic horizontal and vertical motions.

More recently, Chini et al. (2022) have challenged at least some of the consequences of this fundamental concept by noting that the flow field revealed by the DNS of Brethouwer et al. (2007), Augier et al. (2015) and Maffioli and Davidson (2016) does, in fact, exhibit spatio-temporally intermittent motions on small horizontal scales—well-separated from the outer horizontal scale—that are associated with (local) shear instabilities. Using this information, they proposed a new multiscale asymptotic model of stratified turbulence at $Pr = O(1)$, using the concept of ‘marginal stability’ to constrain the representative minimum values of the gradient Richardson number to be $O(1)$, thus determining key properties of ‘fast’ motions associated with the presumed break down of these local shear instabilities. Importantly, this model recovers the usual (and empirically observed) scaling law $H^* \propto FrL^*$ for the vertical eddy scale, but predicts a characteristic vertical velocity scale $W^* \propto Fr^{1/2}U^*$ and a characteristic buoyancy scale $B^* \propto Fr^{3/2}L^*N^{*2}$. These scalings for W^* and B^* both differ from the corresponding predictions of Brethouwer et al. (2007) as a direct consequence of relaxing the assumption of a single horizontal length scale to allow for the idealized modelling of certain important aspects of the spatiotemporally intermittent shear instabilities.

1.2 Stratified turbulence driven by horizontal flows at $Pr \ll 1$

By contrast with geophysical applications, the study of stratified turbulence driven by horizontal flows in stellar interiors is still in its infancy. Most of the early work on the topic is summarized in the seminal paper of Zahn (1992), who estimated the vertical turbulent diffusivity D^* of momentum, or of a passive scalar, in terms of the energy dissipation rate ε^* and of the local fluid properties κ^* and N^* :

$$D^* \propto \left(\frac{\varepsilon^* \kappa^*}{N^{*2}} \right)^{1/2}. \quad (8)$$

To arrive at this conclusion, he made two assumptions: (1) that the vertical diffusivity is primarily due to the largest eddies that are unstable according to (4), where the relevant vertical shear $S^* = u^*(\ell^*)/\ell^*$; and (2) that $\varepsilon^* \propto u^{*3}(\ell^*)/\ell^*$. Lignières (2020) recently revisited Zahn’s argument, interpreting the relevant scale ℓ^* as a “diffusive” or “modified” Ozmidov scale, and explicitly writing it as

$$\ell_{OM}^* = \left(\frac{\kappa^* \varepsilon^{*1/3}}{N^{*2}} \right)^{3/8}. \quad (9)$$

Using critical balance theory, Skoutnev (2023) then argued that it is possible to relate ℓ_{OM}^* and $u^*(\ell_{OM}^*)$ more explicitly to properties of the larger-scale flow (U^* , W^* , H^* and L^*), arriving at the conclusion that

$$H^* \propto \left(\frac{U^* \kappa^*}{N^{*2} L^{*3}} \right)^{1/4} L^* = \left(\frac{Fr^2}{Pe} \right)^{1/4} L^*, \text{ and } W^* \propto \left(\frac{Fr^2}{Pe} \right)^{1/4} U^*, \quad (10a)$$

so

$$D^* \propto H^* W^* \propto \left(\frac{Fr^2}{Pe} \right)^{1/2} U^* L^* = \left(\frac{U^{*3} \kappa^*}{L^* N^{*2}} \right)^{1/2}, \quad (10b)$$

which is consistent with Zahn’s estimate (8) provided one further assumes that $\varepsilon^* \propto U^{*3}/L^*$. Note that this assumption, however, may only be true when the effects of buoyancy are sufficiently weak; see Mashayek et al. (2022).

As turbulence in stars cannot be directly observed, and experimentation with $Pr \ll 1$ fluids is particularly difficult, numerical simulations are best suited to provide insight into the low Pr regime. Cope et al. (2020) and Garaud (2020) recently presented a series of DNS of stratified turbulence driven by horizontal shear at low Pr , focusing on flows where the outer scale Péclet number Pe is low and high, respectively. They found, as in the $Pr = O(1)$ simulations of Brethouwer et al. (2007), that the turbulence becomes highly anisotropic as stratification increases (Fr decreases), and identified three distinct anisotropic regimes: (1) a fully turbulent regime, (2) a regime where the turbulence is spatially and temporally intermittent (as viscosity begins to affect the flow), and (3) a fully viscous regime (where viscosity completely controls the flow dynamics). In the fully turbulent and intermittent regimes, the horizontal motions contain well-separated large and small scales.

When $Pe \ll 1$, diffusion is important at all scales, regardless of the stratification. Cope et al. (2020) found empirically that the vertical size and characteristic velocity of eddies in the fully turbulent regime scale as

$$H^* \propto \left(\frac{Fr^2}{Pe}\right)^{1/3} L^*, \quad W^* \propto \left(\frac{Fr^2}{Pe}\right)^{1/6} U^*, \quad (11)$$

which notably differ from the predictions of Skoutnev (2023). This discrepancy between the DNS results of Cope et al. (2020) and Skoutnev’s theory needs to be explained and is one of the principal motivations for this paper. Interestingly, the prediction for the turbulent diffusivity $D^* \propto H^*W^*$ yields the *same* expression (10b) as in Zahn (1992), Lignières (2020) and Skoutnev (2023), despite H^* and W^* satisfying different scaling laws.

It is important to note, however, that Zahn (1992) did not distinguish between viscous and non-viscous regimes or diffusive and non-diffusive regimes, assuming instead that the turbulence can always develop on scales where diffusion is important but viscous effects are negligible. Yet, the results of Cope et al. (2020) and Garaud (2020) demonstrate that viscosity always eventually begins to affect the turbulence as the stratification increases, so its impact needs to be accounted for in the models. By analogy with Brethouwer et al. (2007) and Chini et al. (2022), one may expect the buoyancy Reynolds number Re_b to be the relevant bifurcation parameter describing the impact of viscosity, a result that will be formally established in this paper. Garaud (2020) also demonstrated numerically the existence of a regime of stratified turbulence at $Pe \gg 1$ and $Pr \ll 1$ where diffusion is negligible. This is not surprising in hindsight, but was not anticipated by Zahn (1992). She tentatively proposed that her data is consistent with $H^* \propto Fr^{2/3}L^*$, and $W^* \propto Fr^{2/3}U^*$ as long as viscosity remains negligible, which is surprising given that one might naively have expected to recover the geophysical regime scalings in that case. She acknowledged, however, that her simulations may not have been performed at a sufficiently high Reynolds number to be in a meaningful asymptotic regime yet. As such, her findings need to be revisited.

The results presented above illustrate the complexity of stratified turbulence at low Pr , and motivate the need for additional work to determine how many regimes exist, how salient flow properties scale with input parameters in each regime, and finally, where the regime boundaries lie in parameter space. In this paper, we therefore approach the problem systematically by adapting the multiscale asymptotic methodology developed by Chini et al. (2022) for the $Pr = O(1)$ regime to the low Pr regime. We begin in § 2 by laying out the model equations and boundary conditions and performing the same anisotropic scaling analysis as Brethouwer et al. (2007), explicitly comparing the $Pr \ll 1$ and the $Pr = O(1)$ cases. This analysis reveals the crucial role of the buoyancy Péclet number (thus named by analogy with Re_b), *viz.* $Pe_b = PrRe_b$. We then perform a slow-fast decomposition of the governing equations in § 3. At high Pe_b , we recover the results of Chini et al. (2022) and highlight the reason why this multiscale approach leads to a different vertical velocity scaling from the one derived by Brethouwer et al. (2007). We then present new results at low Pe_b . Our findings are discussed in § 4, where we use this analysis to partition the parameter space into various regimes of stratified turbulence at low Pr and explicitly provide scaling relationships for the vertical length scale and velocity in each case. Implications and potential applications of our results are presented in § 5.

2 Governing equations and anisotropic scalings

Consider a three-dimensional, non-rotating, incompressible flow expressed in a Cartesian coordinate system where the vertical coordinate z^* (with unit vector \hat{e}_z) is anti-aligned with gravity ($\mathbf{g}^* = -g^*\hat{e}_z$). The fluid is stably stratified, and has a mean density ρ_0^* and a constant background buoyancy frequency N^* . Buoyancy perturbations away from this mean state are incorporated in accordance with the Boussinesq approximation (Spiegel and Veronis, 1960). A body force $F^*\hat{e}_x$ is applied to drive a mean horizontally sheared flow (where F^* is a function of the spanwise variable y^* only, and \hat{e}_x is the unit vector in the streamwise, i.e. x^* , direction). The dimensional governing equations are

$$\frac{\partial \mathbf{u}^*}{\partial t^*} + \mathbf{u}^* \cdot \nabla^* \mathbf{u}^* = -\frac{1}{\rho_0^*} \nabla^* p^* + b^* \hat{e}_z + \nu^* \nabla^{*2} \mathbf{u}^* + \frac{F^*(y^*)}{\rho_0^*} \hat{e}_x, \quad (12a)$$

$$\nabla^* \cdot \mathbf{u}^* = 0, \quad (12b)$$

$$\frac{\partial b^*}{\partial t^*} + \mathbf{u}^* \cdot \nabla^* b^* + N^{*2} w^* = \kappa^* \nabla^{*2} b^*, \quad (12c)$$

where $\nabla^* = (\partial/\partial x^*, \partial/\partial y^*, \partial/\partial z^*)$, $\mathbf{u}^* = (u^*, v^*, w^*)$ denotes the velocity field, p^* the pressure, and b^* the buoyancy perturbation with respect to the background stratification. In accord with the Boussinesq approximation, the fluid has a constant kinematic viscosity ν^* and constant diffusivity κ^* . All perturbations from the stratified rest state are assumed to be triply periodic in a domain of size (L_x^*, L_y^*, L_z^*) . Accordingly, the body force is also assumed to be periodic in y^* , with a characteristic length scale $L^* \leq L_y^*$.

2.1 Anisotropic scalings at $Pr = O(1)$

In the limit of strong stratification, vertical displacements are energetically costly, and hence, as discussed in §1, fluid motions become strongly anisotropic (with $u^*, v^* \gg w^*$). Therefore, it seems natural to non-dimensionalize the governing equations (12) anisotropically. Billant and Chomaz (2001) and Brethouwer et al. (2007) have argued that the (dimensional) horizontal and vertical length scales of turbulent eddies ought to be L^* and H^* , respectively, with an aspect ratio

$$\alpha \equiv \frac{H^*}{L^*} \ll 1, \quad (13)$$

where the dependence of α on the stratification (and other governing parameters) is determined from the following asymptotic analysis. We then introduce a new vertical coordinate ζ^* such that

$$z^* = \alpha \zeta^*. \quad (14)$$

Consequently, if the horizontal velocity scale is U^* , then the vertical velocity scale must be αU^* to respect the divergence free condition without overly restricting the allowable types of flows. Time should be scaled by the turnover time of the horizontal eddies L^*/U^* , and pressure by $\rho_0^* U^{*2}$. To ensure that the nonlinear terms balance the forcing, $U^* = (F_0^* L^*/\rho_0^*)^{1/2}$, where F_0^* is the characteristic forcing amplitude. Finally, the buoyancy scale is chosen to be $H^* N^{*2}$ to ensure that the vertical advection of the background stratification enters the buoyancy equation at leading order and balances the horizontal advection of the buoyancy fluctuations.

Denoting the horizontal components of the velocity as $\mathbf{u}_\perp = (u, v)$, and similarly for the horizontal gradient $\nabla_\perp = (\partial/\partial x, \partial/\partial y)$, the dimensionless system is given by

$$\frac{\partial \mathbf{u}_\perp}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) \mathbf{u}_\perp + w \frac{\partial \mathbf{u}_\perp}{\partial \zeta} = -\nabla_\perp p + \frac{1}{Re\alpha^2} \left(\alpha^2 \nabla_\perp^2 \mathbf{u}_\perp + \frac{\partial^2 \mathbf{u}_\perp}{\partial \zeta^2} \right) + F \hat{e}_x, \quad (15a)$$

$$\frac{\partial w}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) w + w \frac{\partial w}{\partial \zeta} = -\frac{1}{\alpha^2} \frac{\partial p}{\partial \zeta} + \frac{b}{Fr^2} + \frac{1}{Re\alpha^2} \left(\alpha^2 \nabla_\perp^2 w + \frac{\partial^2 w}{\partial \zeta^2} \right), \quad (15b)$$

$$\nabla_\perp \cdot \mathbf{u}_\perp + \frac{\partial w}{\partial \zeta} = 0, \quad (15c)$$

$$\frac{\partial b}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) b + w \frac{\partial b}{\partial \zeta} + w = \frac{1}{Pe\alpha^2} \left(\alpha^2 \nabla_\perp^2 b + \frac{\partial^2 b}{\partial \zeta^2} \right), \quad (15d)$$

where all variables are now non-dimensional and

$$Re = \frac{U^* L^*}{\nu^*}, \quad Fr = \frac{U^*}{N^* L^*}, \quad Pe = \frac{U^* L^*}{\kappa^*} = Pr Re \quad (16)$$

are the usual Reynolds, Froude, and Péclet numbers based on the characteristic horizontal scales of the flow.

When the stratification is strong, $Fr \ll 1$, and the buoyancy term in the vertical component of the momentum equation is unbalanced unless it is compensated by the vertical pressure gradient. In other words, the flow anisotropy implies that hydrostatic balance must be satisfied at lowest order, which then requires

$$\alpha = Fr, \quad (17)$$

implying that the characteristic vertical velocity $W^* = Fr U^*$ while the characteristic vertical scale of the flow $H^* = U^*/N^*$. As discussed in §1, this scaling for H^* is well-established, and has been observed in several laboratory and numerical experiments (Holford and Linden, 1999; Brethouwer et al., 2007; Oglethorpe et al., 2013).

Keeping only the lowest-order terms in an asymptotic expansion of (15) in $\alpha = Fr \ll 1$ yields

$$\frac{\partial \mathbf{u}_\perp}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) \mathbf{u}_\perp + w \frac{\partial \mathbf{u}_\perp}{\partial \zeta} = -\nabla_\perp p + \frac{1}{Re_b} \frac{\partial^2 \mathbf{u}_\perp}{\partial \zeta^2} + F \hat{\mathbf{e}}_x, \quad (18a)$$

$$\frac{\partial p}{\partial \zeta} = b, \quad \nabla_\perp \cdot \mathbf{u}_\perp + \frac{\partial w}{\partial \zeta} = 0, \quad (18b,c)$$

$$\frac{\partial b}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) b + w \frac{\partial b}{\partial \zeta} + w = \frac{1}{Pe_b} \frac{\partial^2 b}{\partial \zeta^2}, \quad (18d)$$

where

$$Re_b = \alpha^2 Re \quad (19)$$

is the usually-defined buoyancy Reynolds number and

$$Pe_b = Pr Re_b = \alpha^2 Pe \quad (20)$$

is the corresponding buoyancy Péclet number. Note that when $Pr = O(1)$, the condition $Re_b \geq O(1)$, which is necessary for viscous effects to be small or negligible, implies that $Pe_b = Pr Re_b \geq O(1)$. As such, the effects of buoyancy diffusion are also small or negligible. The set of equations (18), which will be referred to as the ‘‘anisotropically-scaled high- Pe_b ’’ equations hereafter, recovers the results of Billant and Chomaz (2001) and Brethouwer et al. (2007).

2.2 Anisotropically-scaled equations for $Pr \ll 1$

When the Prandtl number is asymptotically small, however, it is possible to have a regime where $Pe_b \ll 1 \leq Re_b$. In this extreme limit, the diffusion term on the right hand side of (15d) becomes unbalanced, unless the buoyancy field itself is much smaller than anticipated by the scaling $H^* N^{*2}$ used in the previous section. The strongly diffusive limit has in fact already been studied by Lignières (1999), who showed that the buoyancy equation asymptotically reduces to a balance between the vertical advection of the background stratification and the diffusion of the buoyancy fluctuations in that limit. In dimensional terms, we therefore expect $N^{*2} w^* \simeq \kappa^* \nabla^2 b^*$. With this in mind, we let $b = Pe_b \hat{b}$ in (15a)–(15d) (this is equivalent to scaling the dimensional buoyancy by $\alpha^3 N^{*2} U^* L^{*2} / \kappa^*$ instead of $H^* N^{*2}$), and anticipate that $\hat{b} = O(1)$. The resulting dimensionless system becomes

$$\frac{\partial \mathbf{u}_\perp}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) \mathbf{u}_\perp + w \frac{\partial \mathbf{u}_\perp}{\partial \zeta} = -\nabla_\perp p + \frac{1}{Re_b} \left(\alpha^2 \nabla_\perp^2 \mathbf{u}_\perp + \frac{\partial^2 \mathbf{u}_\perp}{\partial \zeta^2} \right) + F \hat{\mathbf{e}}_x, \quad (21a)$$

$$\frac{\partial w}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) w + w \frac{\partial w}{\partial \zeta} = -\frac{1}{\alpha^2} \frac{\partial p}{\partial \zeta} + \frac{Pe_b}{Fr^2} \hat{b} + \frac{1}{Re_b} \left(\alpha^2 \nabla_\perp^2 w + \frac{\partial^2 w}{\partial \zeta^2} \right), \quad (21b)$$

$$\nabla_\perp \cdot \mathbf{u}_\perp + \frac{\partial w}{\partial \zeta} = 0, \quad (21c)$$

$$\frac{\partial \hat{b}}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) \hat{b} + w \frac{\partial \hat{b}}{\partial \zeta} + \frac{1}{Pe_b} w = \frac{1}{Pe_b} \left(\alpha^2 \nabla_\perp^2 \hat{b} + \frac{\partial^2 \hat{b}}{\partial \zeta^2} \right). \quad (21d)$$

For sufficiently strong stratification, the vertical component of the momentum equation must again be in hydrostatic balance, which requires

$$\alpha^2 = \frac{Fr^2}{Pe_b} = \frac{Fr^2}{\alpha^2 Pe} \rightarrow \alpha = Fr_M, \text{ where } Fr_M = \left(\frac{Fr^2}{Pe} \right)^{1/4} \quad (22)$$

is a modified Froude number (see Lignières, 2020; Skoutnev, 2023). We therefore find that the characteristic vertical length scale should be $H^* = (Fr^2/Pe)^{1/4} L^*$, and the characteristic vertical velocity scale should be $W^* = (Fr^2/Pe)^{1/4} U^*$, recovering the results of Skoutnev (2023), albeit using a different argument.

In the limit $Fr_M \ll 1$ and $Pe_b \ll 1$, keeping only the lowest order terms in (21a)–(21d) yields

$$\frac{\partial \mathbf{u}_\perp}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) \mathbf{u}_\perp + w \frac{\partial \mathbf{u}_\perp}{\partial \zeta} = -\nabla_\perp p + \frac{1}{Re_b} \frac{\partial^2 \mathbf{u}_\perp}{\partial \zeta^2} + F \hat{\mathbf{e}}_x, \quad (23a)$$

$$\frac{\partial p}{\partial \zeta} = \hat{b}, \quad \nabla_\perp \cdot \mathbf{u}_\perp + \frac{\partial w}{\partial \zeta} = 0, \quad w = \frac{\partial^2 \hat{b}}{\partial \zeta^2}. \quad (23b,c,d)$$

These scaling laws and governing equations are the low Pe_b analogs of equations (18). In what follows, we therefore refer to them as the “anisotropically-scaled low- Pe_b ” equations.

2.3 Evidence for multiscale dynamics

The two sets of anisotropically-scaled equations given by (18) for $Pe_b \geq O(1)$ and (23) for $Pe_b \ll 1$ assume, by construction, that horizontal scales are large, while the vertical scale is small. As such, they necessarily describe the dynamics of weakly-coupled “pancake” vortices or horizontal meanders of the mean flow. They cannot, however, capture the small-scale turbulence that is expected to develop from shear instabilities between these layerwise horizontal motions (Chini et al., 2022). Yet, these instabilities are ubiquitous when Re_b is large enough, and have been observed in laboratory experiments, inferred from oceanographic *in situ* measurements, as well as in direct numerical simulations at $Pr \gtrsim O(1)$ (Gregg et al., 2018; Caulfield, 2021).

Furthermore, recent DNS of stratified turbulence in the $Pe \ll 1$ regime by Cope et al. (2020) also reveal that flows on small horizontal scales are important. As discussed earlier, they empirically find that $H^* \propto (Fr^2/Pe)^{1/3} L^*$ instead of $H^* \propto (Fr^2/Pe)^{1/4} L^*$ and that $W^* \propto (Fr^2/Pe)^{1/6} U^*$ instead of $W^* \propto (Fr^2/Pe)^{1/4} U^*$. These results thus shed some doubt on the relevance of the low- Pe_b anisotropically-scaled equations (23) for modeling stratified turbulence, at least at large buoyancy Reynolds number.

As discussed in § 1, Chini et al. (2022) have recently argued in the context of geophysical stratified turbulence that one *must* take into account the fast, small horizontal scales and study their (marginally-stable) interaction with the slow, larger-scale anisotropic flow to obtain a more complete and more accurate model of stratified turbulence. Accordingly, we now propose to extend their work to the low Pr limit. The next section first outlines the work of Chini et al. (2022) for pedagogical clarity, then extends the analysis to the low Pr regime.

3 Multiscale models for stratified turbulence

We consider the same model set-up as introduced in §2. Here, however, we make no assumption about the amplitude of the vertical flow motions when non-dimensionalising the governing equations, and instead allow it to emerge naturally from the analysis. Accordingly, we non-dimensionalize the vertical velocity by U^* , and correspondingly choose the buoyancy scale to be $L^* N^{*2}$. Then, the dimensionless system is

$$\frac{\partial \mathbf{u}_\perp}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) \mathbf{u}_\perp + \frac{w}{\alpha} \frac{\partial \mathbf{u}_\perp}{\partial \zeta} = -\nabla_\perp p + \frac{1}{Re_b} \left(\alpha^2 \nabla_\perp^2 \mathbf{u}_\perp + \frac{\partial^2 \mathbf{u}_\perp}{\partial \zeta^2} \right) + F \hat{\mathbf{e}}_x, \quad (24a)$$

$$\frac{\partial w}{\partial t} + (\mathbf{u}_\perp \cdot \nabla_\perp) w + \frac{w}{\alpha} \frac{\partial w}{\partial \zeta} = -\frac{1}{\alpha} \frac{\partial p}{\partial \zeta} + \frac{b}{Fr^2} + \frac{1}{Re_b} \left(\alpha^2 \nabla_\perp^2 w + \frac{\partial^2 w}{\partial \zeta^2} \right), \quad (24b)$$

$$\nabla_{\perp} \cdot \mathbf{u}_{\perp} + \frac{1}{\alpha} \frac{\partial w}{\partial \zeta} = 0, \quad (24c)$$

$$\frac{\partial b}{\partial t} + (\mathbf{u}_{\perp} \cdot \nabla_{\perp}) b + \frac{w}{\alpha} \frac{\partial b}{\partial \zeta} + w = \frac{1}{Pe_b} \left(\alpha^2 \nabla_{\perp}^2 b + \frac{\partial^2 b}{\partial \zeta^2} \right). \quad (24d)$$

These equations are the starting point for our analysis. We assume that $Re_b \geq O(1)$, and that Fr is sufficiently small to ensure that $\alpha \ll 1$, but make no other *a priori* assumption on the size of Pe_b at this stage.

3.1 Slow-fast decomposition

We now perform a multiscale expansion of the system in the limit of small aspect ratio α . Following Chini et al. (2022), we assume the existence of two distinct horizontal length scales: the original large scales that are $O(1)$ in the chosen nondimensionalization, as well as much smaller horizontal scales that are $O(\alpha)$. With that choice, small-scale fluid motions are isotropic by construction. We further assume that the flow has two distinct time scales: a slow time scale associated with the turnover of the large horizontal eddies, as before, and a fast time scale inversely related to the vertical shearing rate of the large-scale mean flow U^*/H^* . In practice, we thus define the slow and fast horizontal coordinates as $\mathbf{x}_s = \mathbf{x}_{\perp}$ and $\mathbf{x}_f = \mathbf{x}_s/\alpha$, respectively (henceforth, the subscript f denotes fast and s denotes slow). Similarly, we split time into slow and fast variables, such that $t_f = t_s/\alpha$ where $t_s = t$. Consequently, the partial derivatives with respect to time and to the horizontal variables become

$$\frac{\partial}{\partial t} = \frac{1}{\alpha} \frac{\partial}{\partial t_f} + \frac{\partial}{\partial t_s}, \quad \nabla_{\perp} = \frac{1}{\alpha} \nabla_f + \nabla_s. \quad (25)$$

All dependent variables (collectively referred to as q) are now assumed to be functions of both fast and slow length and time scales: $q = q(\mathbf{x}_f, \mathbf{x}_s, \zeta, t_f, t_s; \alpha)$.

Assuming the fast and slow scales are sufficiently separated, Chini et al. (2022) then define a fast-averaging operator $\overline{(\cdot)}$, such that

$$\overline{q}(\mathbf{x}_s, \zeta, t_s; \alpha) = \lim_{T, l_x, l_y \rightarrow \infty} \frac{1}{l_x l_y T} \int_0^T \int_{\mathcal{D}} q(\mathbf{x}_f, \mathbf{x}_s, \zeta, t_f, t_s; \alpha) d\mathbf{x}_f dt_f, \quad (26)$$

where \mathcal{D} is a horizontal domain, with fast spatial periods l_x and l_y , and T is the fast time-integration period. With this definition, \overline{q} depends on slow variables only. Each quantity q can then be split into a slowly-varying field \overline{q} and a rapidly fluctuating component $q' = q - \overline{q}$, which implies that the fast-average of the fluctuation field must vanish, i.e., $\overline{q'} = 0$. Note that q' itself could, however, still depend on the slow length and time scales.

We first substitute the expressions (25) for ∇_{\perp} and $\partial/\partial t$ into (24), and split each field q into $\overline{q} + q'$. We then take the fast average of each of the four governing equations to obtain the evolution equations for the mean flow, then subtract the mean from the total to obtain the evolution equations for the fluctuations.

Starting with the continuity equation, we have

$$\frac{1}{\alpha} \nabla_f \cdot \mathbf{u}'_{\perp} + \nabla_s \cdot \overline{\mathbf{u}}_{\perp} + \nabla_s \cdot \mathbf{u}'_{\perp} + \frac{1}{\alpha} \frac{\partial \overline{w}}{\partial \zeta} + \frac{1}{\alpha} \frac{\partial w'}{\partial \zeta} = 0, \quad (27)$$

whose fast-average reveals that

$$\nabla_s \cdot \overline{\mathbf{u}}_{\perp} + \frac{1}{\alpha} \frac{\partial \overline{w}}{\partial \zeta} = 0 \quad (28a)$$

for the mean flow and

$$\frac{1}{\alpha} \nabla_f \cdot \mathbf{u}'_{\perp} + \nabla_s \cdot \mathbf{u}'_{\perp} + \frac{1}{\alpha} \frac{\partial w'}{\partial \zeta} = 0 \quad (28b)$$

for the perturbations.

A similar procedure for the horizontal momentum equation yields

$$\begin{aligned} \frac{\partial \bar{\mathbf{u}}_{\perp}}{\partial t_s} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_s \bar{\mathbf{u}}_{\perp} + \frac{\bar{w}}{\alpha} \frac{\partial \bar{\mathbf{u}}_{\perp}}{\partial \zeta} + \frac{1}{\alpha} \left(\overline{\mathbf{u}'_{\perp} \cdot \nabla_f \mathbf{u}'_{\perp} + w' \frac{\partial \mathbf{u}'_{\perp}}{\partial \zeta}} \right) + \overline{\mathbf{u}'_{\perp} \cdot \nabla_s \mathbf{u}'_{\perp}} \\ = -\nabla_s \bar{p} + \frac{1}{Re_b} \left(\frac{\partial^2 \bar{\mathbf{u}}_{\perp}}{\partial \zeta^2} + \alpha^2 \nabla_s^2 \bar{\mathbf{u}}_{\perp} \right) + \bar{F}, \end{aligned} \quad (29a)$$

for the mean flow (where $F = \bar{F} = O(1)$ by construction), and

$$\begin{aligned} \frac{1}{\alpha} \frac{\partial \mathbf{u}'_{\perp}}{\partial t_f} + \frac{\partial \mathbf{u}'_{\perp}}{\partial t_s} + \frac{1}{\alpha} \bar{\mathbf{u}}_{\perp} \cdot \nabla_f \mathbf{u}'_{\perp} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_s \mathbf{u}'_{\perp} + \mathbf{u}'_{\perp} \cdot \nabla_s \bar{\mathbf{u}}_{\perp} + \frac{1}{\alpha} \mathbf{u}'_{\perp} \cdot \nabla_f \mathbf{u}'_{\perp} + \mathbf{u}'_{\perp} \cdot \nabla_s \mathbf{u}'_{\perp} \\ + \frac{1}{\alpha} \left(\bar{w} \frac{\partial \mathbf{u}'_{\perp}}{\partial \zeta} + w' \frac{\partial \bar{\mathbf{u}}_{\perp}}{\partial \zeta} + w' \frac{\partial \mathbf{u}'_{\perp}}{\partial \zeta} \right) = -\frac{1}{\alpha} \nabla_f p' + \frac{1}{Re_b} \left(\nabla_f^2 \mathbf{u}'_{\perp} + \frac{\partial^2 \mathbf{u}'_{\perp}}{\partial \zeta^2} \right) \\ + \frac{1}{\alpha} \left(\overline{\mathbf{u}'_{\perp} \cdot \nabla_f \mathbf{u}'_{\perp} + w' \frac{\partial \mathbf{u}'_{\perp}}{\partial \zeta}} \right) + \overline{\mathbf{u}'_{\perp} \cdot \nabla_s \mathbf{u}'_{\perp}}, \end{aligned} \quad (29b)$$

for the fluctuations.

The mean buoyancy equation is

$$\frac{\partial \bar{b}}{\partial t_s} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_s \bar{b} + \frac{\bar{w}}{\alpha} \frac{\partial \bar{b}}{\partial \zeta} + \frac{1}{\alpha} \left(\overline{\mathbf{u}'_{\perp} \cdot \nabla_f b' + w' \frac{\partial \bar{b}}{\partial \zeta}} \right) + \overline{\mathbf{u}'_{\perp} \cdot \nabla_s \bar{b}} + \bar{w} = \frac{1}{Pe_b} \left(\frac{\partial^2 \bar{b}}{\partial \zeta^2} + \alpha^2 \nabla_s^2 \bar{b} \right), \quad (30a)$$

while the corresponding fluctuation equation becomes:

$$\begin{aligned} \frac{1}{\alpha} \frac{\partial b'}{\partial t_f} + \frac{\partial b'}{\partial t_s} + \frac{1}{\alpha} \bar{\mathbf{u}}_{\perp} \cdot \nabla_f b' + \mathbf{u}'_{\perp} \cdot \nabla_s \bar{b} + \frac{1}{\alpha} \mathbf{u}'_{\perp} \cdot \nabla_f b' + \mathbf{u}'_{\perp} \cdot \nabla_s b' \\ + \frac{1}{\alpha} \left(w' \frac{\partial \bar{b}}{\partial \zeta} + \bar{w} \frac{\partial b'}{\partial \zeta} + w' \frac{\partial b'}{\partial \zeta} \right) + w' = \frac{1}{Pe_b} \left(\nabla_f^2 b' + \frac{\partial^2 b'}{\partial \zeta^2} \right) \\ + \frac{1}{\alpha} \left(\overline{\mathbf{u}'_{\perp} \cdot \nabla_f b' + w' \frac{\partial b'}{\partial \zeta}} \right) + \overline{\mathbf{u}'_{\perp} \cdot \nabla_s b'}. \end{aligned} \quad (30b)$$

Finally, the mean vertical momentum equation is

$$\begin{aligned} \frac{\partial \bar{w}}{\partial t_s} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_s \bar{w} + \frac{\bar{w}}{\alpha} \frac{\partial \bar{w}}{\partial \zeta} + \frac{1}{\alpha} \left(\overline{\mathbf{u}'_{\perp} \cdot \nabla_f w' + w' \frac{\partial w'}{\partial \zeta}} \right) + \overline{\mathbf{u}'_{\perp} \cdot \nabla_s w'} \\ = -\frac{1}{\alpha} \frac{\partial \bar{p}}{\partial \zeta} + \frac{\bar{b}}{Fr^2} + \frac{1}{Re_b} \left(\frac{\partial^2 \bar{w}}{\partial \zeta^2} + \alpha^2 \nabla_s^2 \bar{w} \right), \end{aligned} \quad (31a)$$

while the fluctuations satisfy

$$\begin{aligned} \frac{1}{\alpha} \frac{\partial w'}{\partial t_f} + \frac{\partial w'}{\partial t_s} + \frac{1}{\alpha} \bar{\mathbf{u}}_{\perp} \cdot \nabla_f w' + \mathbf{u}'_{\perp} \cdot \nabla_s \bar{w} + \frac{1}{\alpha} \mathbf{u}'_{\perp} \cdot \nabla_f w' + \mathbf{u}'_{\perp} \cdot \nabla_s w' \\ + \frac{1}{\alpha} \left(w' \frac{\partial \bar{w}}{\partial \zeta} + \bar{w} \frac{\partial w'}{\partial \zeta} + w' \frac{\partial w'}{\partial \zeta} \right) = -\frac{1}{\alpha} \frac{\partial p'}{\partial \zeta} + \frac{b'}{Fr^2} + \frac{1}{Re_b} \left(\nabla_f^2 w' + \frac{\partial^2 w'}{\partial \zeta^2} \right) \\ + \frac{1}{\alpha} \left(\overline{\mathbf{u}'_{\perp} \cdot \nabla_f w' + w' \frac{\partial w'}{\partial \zeta}} \right) + \overline{\mathbf{u}'_{\perp} \cdot \nabla_s w'}. \end{aligned} \quad (31b)$$

We see, as noted by Chini et al. (2022), that the effective Reynolds and Péclet numbers of the fluctuation equations are Re_b/α and Pe_b/α , respectively, which implies that the fluctuations are formally much *less* viscous and *less* diffusive than the mean. This perhaps counterintuitive conclusion is a direct consequence of the flow anisotropy.

3.2 Multiscale model at $Pe_b \geq O(1)$

We begin by summarizing the steps taken by Chini et al. (2022) to derive a reduced multiscale model for stratified turbulence at $Re_b, Pe_b \geq O(1)$, as much of the analysis proves to be similar at low Prandtl number. Following that work, we posit the following asymptotic expansions:

$$[b, p, \mathbf{u}_\perp, w] \sim [b_0, p_0, \mathbf{u}_{\perp 0}, w_0] + \alpha^{1/2}[b_1, p_1, \mathbf{u}_{\perp 1}, w_1] + \alpha[b_2, p_2, \mathbf{u}_{\perp 2}, w_2] + \dots \quad (32)$$

The expansions start at $O(1)$ to reflect the expectation that the dominant contributions to the pressure and the horizontal velocity arise on large horizontal scales. The expansions for b and w also start at $O(1)$, although we readily show below that w_0 and b_0 must both vanish when $\alpha \rightarrow 0$. Finally, the expansions proceed as asymptotic series in $\alpha^{1/2}$ following the results of Chini et al. (2022). They demonstrated that, because the small-scale fluctuations are isotropic, \mathbf{u}' and w' are necessarily of the same order. Inspection of the mean horizontal momentum equation then immediately reveals that both fields need to be $O(\alpha^{1/2})$ to ensure that the Reynolds stresses feedback on $\mathbf{u}_{\perp 0}$ at leading order. (See below for further details.)

With these choices, we substitute (32) into the equations with multiscale derivatives presented above, analysing in turn the continuity equation (28), the horizontal component of the momentum equation (29), the buoyancy equation (30), and finally, the vertical component of the momentum equation (31). At each step, we match terms at leading order to infer their sizes and respective evolution equations, and thus derive a reduced model for the flow.

Considering first the mean continuity equation (28a), we see that $\partial_\zeta \bar{w}_0 = \partial_\zeta \bar{w}_1 = 0$, implying $\bar{w}_0 = \bar{w}_1 = 0$ to suppress unphysical ‘elevator modes’ (that are allowed by the vertically periodic boundary conditions) from our model. Consequently,

$$\nabla_s \cdot \bar{\mathbf{u}}_{\perp 0} + \frac{\partial \bar{w}_2}{\partial \zeta} = 0. \quad (33a)$$

In the fluctuation continuity equation (28b), the second term is clearly much smaller than the first and can therefore be neglected from the leading-order set of dominant terms. Substituting the asymptotic series (32) we then see that

$$\nabla_f \cdot \mathbf{u}'_{\perp i} + \frac{\partial w'_i}{\partial \zeta} = 0, \quad (33b)$$

for $i = 0, 1$ (for larger values of i , the slow derivative of $\mathbf{u}'_{\perp i-2}$ should be taken into account).

We now turn to the mean horizontal component of the momentum equation (29a). The Reynolds stresses must be $O(1)$ to feed back on the mean flow, so \mathbf{u}' and w' must both be $O(\alpha^{1/2})$. Hence, $\mathbf{u}'_{\perp 0} = \mathbf{0}$ and $w'_0 = 0$, yielding $\mathbf{u}_{\perp 0} = \bar{\mathbf{u}}_{\perp 0}$ and that $w_0 = 0$, since $\bar{w}_0 = 0$, too. (See Chini et al., 2022, for a more detailed discussion of why $\mathbf{u}'_{\perp 0} = \mathbf{0}$.) In the horizontal momentum equation for the fluctuations, the fast dynamics take place at $O(\alpha^{-1/2})$ since $\mathbf{u}' = O(\alpha^{1/2})$. Ensuring that pressure is a leading-order effect implies that $p'_0 = 0$, so $p_0 = \bar{p}_0$, as expected. Many of the remaining terms are formally higher order, including all fluctuation-fluctuation interactions, which are $O(1)$ or smaller. Of the nonlinear terms, the only ones that contribute at leading order are quasilinear: $\alpha^{-1} \bar{\mathbf{u}}_\perp \cdot \nabla_f \mathbf{u}'_\perp$ and $\alpha^{-1} w' \partial_z \bar{\mathbf{u}}_\perp$. Therefore, after substituting (32) into (29) and retaining only the leading terms, we obtain

$$\frac{\partial \bar{\mathbf{u}}_{\perp 0}}{\partial t_s} + \bar{\mathbf{u}}_{\perp 0} \cdot \nabla_s \bar{\mathbf{u}}_{\perp 0} + \bar{w}_2 \frac{\partial \bar{\mathbf{u}}_{\perp 0}}{\partial \zeta} = -\nabla_s \bar{p}_0 - \frac{\partial}{\partial \zeta} \left(\overline{w'_1 \mathbf{u}'_{\perp 1}} \right) + \frac{1}{Re_b} \frac{\partial^2 \bar{\mathbf{u}}_{\perp 0}}{\partial \zeta^2} + \bar{F}, \quad (34a)$$

$$\frac{\partial \mathbf{u}'_{\perp 1}}{\partial t_f} + \bar{\mathbf{u}}_{\perp 0} \cdot \nabla_f \mathbf{u}'_{\perp 1} + w'_1 \frac{\partial \bar{\mathbf{u}}_{\perp 0}}{\partial \zeta} = -\nabla_f p'_1 + \frac{\alpha}{Re_b} \left(\nabla_f^2 \mathbf{u}'_{\perp 1} + \frac{\partial^2 \mathbf{u}'_{\perp 1}}{\partial \zeta^2} \right), \quad (34b)$$

where the formally higher-order Laplacian term has been retained to regularize the fluctuation equation.

Next, we examine the buoyancy equation, which reveals the sizes of \bar{b} and b' . In the mean equation (30a), $\bar{w} = O(\alpha)$, and the buoyancy flux term is formally also of that order (see Chini et al., 2022, and also below). Thus, $\bar{b} = O(\alpha)$ as well, as long as $Pe_b \geq O(1)$, which is implicit in the regime considered in this section, implying $\bar{b}_0 = \bar{b}_1 = 0$. The size of the buoyancy flux can be confirmed by inspection of the buoyancy fluctuation equation (30b). To be in a regime in which the stratification impacts the turbulent motions, the $O(b'/\alpha)$ fast dynamics must be of the same order as the advection of the background stratification, which is $O(w') = O(\alpha^{1/2})$. This ordering implies that $b' = O(\alpha^{3/2})$, so $b'_0 = b'_1 = b'_2 = 0$. Consequently, $b_0 = b_1 = 0$, and $b_2 = \bar{b}_2$. Fluctuation-fluctuation interactions in this equation are again formally higher order, and the

only remaining nonlinearities are quasilinear. At leading order, using (32) in (30) shows that the mean and fluctuation buoyancy equations are

$$\frac{\partial \bar{b}_2}{\partial t_s} + \bar{\mathbf{u}}_{\perp 0} \cdot \nabla_s \bar{b}_2 + \bar{w}_2 \frac{\partial \bar{b}_2}{\partial \zeta} + \bar{w}_2 = -\frac{\partial}{\partial \zeta} \left(\overline{w'_1 b'_3} \right) + \frac{1}{Pe_b} \frac{\partial^2 \bar{b}_2}{\partial \zeta^2}, \quad (35a)$$

$$\frac{\partial b'_3}{\partial t_f} + \bar{\mathbf{u}}_{\perp 0} \cdot \nabla_f b'_3 + w'_1 \frac{\partial \bar{b}_2}{\partial \zeta} + w'_1 = \frac{\alpha}{Pe_b} \left(\nabla_f^2 b'_3 + \frac{\partial^2 b'_3}{\partial \zeta^2} \right), \quad (35b)$$

where as before the diffusion term in the fluctuation equation has been retained for regularization.

Finally, we identify the leading order terms in the vertical component of the momentum equation. In the mean equation (31a), a hydrostatic leading-order balance requires $\alpha = Fr$, as in the anisotropically-scaled model described in § 2.1. This specification, when applied to the fluctuation equation (31b), is quite satisfactory as it implies that the fluctuating buoyancy force b'/Fr^2 arises at leading order. It can also easily be shown that, once again, fluctuation-fluctuation interactions are negligible at leading order, so the fast dynamics are quasilinear. Substituting (32) into (31) and using all of the information available, we obtain the leading order mean and fluctuating components of the vertical momentum equation,

$$\frac{\partial \bar{p}_0}{\partial \zeta} = \bar{b}_2, \quad (36a)$$

$$\frac{\partial w'_1}{\partial t_f} + \bar{\mathbf{u}}_{\perp 0} \cdot \nabla_f w'_1 = -\frac{\partial p'_1}{\partial \zeta} + b'_3 + \frac{\alpha}{Re_b} \left(\nabla_f^2 w'_1 + \frac{\partial^2 w'_1}{\partial \zeta^2} \right), \quad (36b)$$

where the formally higher-order viscous term has been retained to regularize the fluctuation equation.

The system of equations formed by (33) (with $i = 1$), (34), (35) and (36), is equivalent to equations (2.28)–(2.35) in Chini et al. (2022), the only differences arising from a different choice of non-dimensionalization. These equations are fully closed and therefore self-consistent. Crucially, the resulting system is quasilinear, and can be solved through appealing to (plausible and empirically supported) marginal stability arguments for the fluctuations as discussed by Michel and Chini (2019) and Chini et al. (2022).

These equations are valid whenever $\alpha \ll 1$ (equivalently, $Fr \ll 1$ since $\alpha = Fr$), and the buoyancy Reynolds and Péclet numbers are both $O(1)$ or larger, thus allowing for the growth and saturation of the fluctuations via interactions with the mean. When $Pr = O(1)$, $Re_b \geq O(1)$ necessarily implies $Pe_b \geq O(1)$, so the two conditions are equivalent. Since $Re_b = \alpha^2 Re$, this condition is equivalent to $Re, Pe \geq O(Fr^{-2})$.

For low Pr , these reduced equations remain valid as long as $Re_b, Pe_b \geq O(1)$. However, for *sufficiently* low Pr it is possible to have an intermediate regime where $Re_b \geq O(1)$ while $Pe_b = Pr Re_b \ll 1$, and that regime is *not* captured by the reduced equations derived here and in Chini et al. (2022). Yet, this scenario is likely to be relevant in stellar interiors, where Pr is asymptotically small (Garaud et al., 2015b). We now perform a similar analysis to develop a multiscale model that is valid for low- Pe_b flows.

3.3 Multiscale model for $Pe_b \ll 1$

In the limit of small Pe_b , the arguments presented above continue to apply for the continuity equation and the horizontal component of the momentum equation, neither of which involve buoyancy terms. However, the procedure subsequently fails because diffusive effects dominate in the buoyancy equation and are unbalanced unless b is much smaller than expected, mirroring the argument made in § 2.2. To capture this correctly within the context of our proposed multiscale model, we must introduce a two-parameter expansion for small α and small Pe_b in lieu of (32).

As in Chini et al. (2022), we assume that the asymptotic expansion in α proceed in powers of $\alpha^{1/2}$ and show the self-consistency of this choice below. Inspired by Lignières (1999), we now also assume that each field can be expanded as a series in powers of Pe_b as well. We therefore have

$$q = q_{00} + \alpha^{1/2} q_{01} + \alpha q_{02} + \dots + Pe_b \left(q_{10} + \alpha^{1/2} q_{11} + \alpha q_{12} + \dots \right) + O(Pe_b^2), \quad (37)$$

for $q \in \{\mathbf{u}_{\perp}, p, w, b\}$, and assume that $\bar{\mathbf{u}}_{\perp 00} = O(1)$ and $\bar{p}_{00} = O(1)$ to balance the forcing. We then proceed exactly as before, substituting (37) in turn into the multiscale continuity equation (28), the horizontal

component of the momentum equation (29), the buoyancy equation (30), and the vertical component of the momentum equation (31), to extract the relevant reduced equations at leading order.

Starting with the mean continuity equation (28a), we find again that $\bar{w} = O(\alpha)$ and hence $\bar{w}_{00} = \bar{w}_{01} = 0$. Equation (28b) further implies that $O(\mathbf{u}'_{\perp}) = O(w')$, so

$$\nabla_s \cdot \bar{\mathbf{u}}_{\perp 00} + \frac{\partial \bar{w}_{02}}{\partial \zeta} = 0, \quad (38a)$$

$$\nabla_f \cdot \mathbf{u}'_{\perp 01} + \frac{\partial w'_{01}}{\partial \zeta} = 0. \quad (38b)$$

From the mean horizontal component of the momentum equation (29a), \mathbf{u}'_{\perp} and w' must both be $O(\alpha^{1/2})$ as before, hence $\mathbf{u}'_{\perp 00} = 0$ and $w'_{00} = 0$, so $\mathbf{u}_{\perp 00} = \bar{\mathbf{u}}_{\perp 00}$ and $w_{00} = 0$. The inferred sizes of the velocity fluctuations again shows that fluctuation-fluctuation interactions are formally higher-order, and the only remaining nonlinearities in the corresponding fluctuation equation (29b) are quasilinear. Finally, for the fluctuating horizontal pressure gradient to influence the fluctuations of horizontal velocity at leading order, $p' = O(\mathbf{u}'_{\perp}) = O(\alpha^{1/2})$, hence $p'_{00} = 0$. Substituting (37) into (29a) and (29b), using these deductions and retaining only the lowest-order terms, shows that

$$\frac{\partial \bar{\mathbf{u}}_{\perp 00}}{\partial t_s} + (\bar{\mathbf{u}}_{\perp 00} \cdot \nabla_s) \bar{\mathbf{u}}_{\perp 00} + \bar{w}_{02} \frac{\partial \bar{\mathbf{u}}_{\perp 00}}{\partial \zeta} = -\nabla_s \bar{p}_{00} - \frac{\partial}{\partial \zeta} \left(\overline{w'_{01} \mathbf{u}'_{\perp 01}} \right) + \frac{1}{Re_b} \frac{\partial^2 \bar{\mathbf{u}}_{\perp 00}}{\partial \zeta^2} + F \hat{\mathbf{e}}_x, \quad (39a)$$

$$\frac{\partial \mathbf{u}'_{\perp 01}}{\partial t_f} + (\bar{\mathbf{u}}_{\perp 00} \cdot \nabla_f) \mathbf{u}'_{\perp 01} + w'_{01} \frac{\partial \bar{\mathbf{u}}_{\perp 00}}{\partial \zeta} = -\nabla_f p'_{01} + \frac{\alpha}{Re_b} \left(\nabla_f^2 \mathbf{u}'_{\perp 01} + \frac{\partial^2 \mathbf{u}'_{\perp 01}}{\partial \zeta^2} \right), \quad (39b)$$

where the viscous term in the fluctuation equation has been retained for regularization.

Thus far, each step in the analysis has been identical to that taken in the previous section. Rapid diffusion, however, affects the size of the mean and fluctuating buoyancy fields \bar{b} and b' , and does so in different ways because the effective Péclet number of the fluctuations is larger than that of the mean flow. More specifically, having assumed in this section that $Pe_b \ll 1$, we see that two possibilities arise when $\alpha \ll 1$: either $\alpha \ll Pe_b \ll 1$, in which case diffusion is dominant in the mean buoyancy equation but negligible in the fluctuation buoyancy equation, or $Pe_b \ll \alpha$ in which case diffusion is dominant at all scales. In what follows, we investigate both cases in turn.

Before doing so, however, we note that the mean buoyancy equation (30a) in both cases is unbalanced unless $\bar{b} = O(\alpha Pe_b)$ to match the \bar{w} term (again mirroring the arguments given in § 2.2). This implies $\bar{b}_{0i} = 0, \forall i$, and $\bar{b}_{10} = \bar{b}_{11} = 0$. At lowest order in Pe_b , the only surviving terms in (30a) are therefore

$$\frac{1}{\alpha} \left(\overline{\mathbf{u}'_{\perp} \cdot \nabla_f b' + w' \frac{\partial b'}{\partial \zeta}} \right) + \bar{w} = \frac{1}{Pe_b} \frac{\partial^2 \bar{b}}{\partial \zeta^2}, \quad (40)$$

where the size of b' is yet to be determined and differs depending on the relative sizes of Pe_b and α . For this reason, we have retained the turbulent buoyancy flux for now.

3.3.1 Case 1: $\alpha \ll Pe_b \ll 1$ (the intermediate regime)

We first consider a scenario in which $\alpha \ll Pe_b \ll 1$ and henceforth refer to this part of parameter space as the ‘‘intermediate regime’’. While diffusion dominates the mean buoyancy equation, the fact that $\alpha/Pe_b \ll 1$ implies that it only formally enters the buoyancy fluctuation equation (30b) at higher order. Because of this, the evolution of b' is very similar to that obtained in Chini et al. (2022). As in § 3.2, we ensure that the background stratification influences the fast dynamics of b' by requiring $b'/\alpha = O(w') = O(\alpha^{1/2})$, so $b' = O(\alpha^{3/2})$. This implies that $b'_{00}, b'_{01}, b'_{02} = 0$, but $b'_{03} \neq 0$.

Substituting the ansatz (37) into the mean and fluctuation buoyancy equations and using the information collected so far we therefore have

$$\frac{\partial}{\partial \zeta} \left(\overline{w'_{01} b'_{03}} \right) + \bar{w}_{02} = \frac{\partial^2 \bar{b}_{12}}{\partial \zeta^2}, \quad (41a)$$

$$\frac{\partial b'_{03}}{\partial t_f} + \bar{\mathbf{u}}_{\perp 00} \cdot \nabla_f b'_{03} + w'_{01} = \frac{\alpha}{Pe_b} \left(\nabla_f^2 b'_{03} + \frac{\partial^2 b'_{03}}{\partial \zeta^2} \right), \quad (41b)$$

where the diffusion term for the fluctuations can be retained to regularize the equation, but is formally higher order.

We note that the reduced buoyancy fluctuation equation in this regime differs slightly from the one derived by Chini et al. (2022) given in (35b), because it does not contain a term of the form $w'_{01} \partial \bar{b} / \partial \zeta$, which is formally higher-order when $Pe_b \ll 1$. Also, we see that the mean buoyancy equation in that regime differs from the asymptotic low Péclet number (LPN) equations of Lignières (1999), which would not contain a turbulent flux term. This discrepancy arises because his derivation assumes that all dynamics are diffusive, whereas in this intermediate regime the fluctuation dynamics are not and can therefore influence the mean buoyancy field at leading order.

Finally, we examine the vertical component of the momentum equation. Based on past experience in the $Pe_b \geq O(1)$ case (see § 3.2), one would naively expect to recover hydrostatic equilibrium at leading order in the mean vertical momentum equation (31a). Because $\bar{b} = O(\alpha Pe_b)$, this would imply $\alpha = (Fr^2/Pe_b)^{1/2}$ as in the low- Pe_b anisotropically-scaled equations of § 2.2. However, that choice leads to an irreconcilable inconsistency in the fluctuation equation (31b): the fluctuation pressure gradient term is $O(\alpha^{-1/2})$, as are the fast inertial dynamics of w' , but the fluctuation buoyancy term is $O(\alpha^{3/2}/Fr^2) = O(\alpha^{-1/2}/Pe_b)$, which is formally much larger than any other term and is therefore unbalanced. In other words, we cannot reconcile hydrostatic equilibrium at leading order for the mean flow with a balanced equation for the fluctuation w' .

The solution to this conundrum is to insist instead that the equation for w' be balanced, in which case $O(\alpha^{-1/2}) = O(\alpha^{3/2} Fr^{-2})$, thereby recovering the standard scaling relationship $\alpha = Fr$ (Billant and Chomaz, 2001; Brethouwer et al., 2007; Chini et al., 2022). With this choice, the leading-order vertical pressure gradient in the mean equation is asymptotically small (e.g. as for the wall-normal pressure gradient in laminar boundary-layer theory of Batchelor, 1967). This perhaps unexpected result is discussed in § 4 below. Substituting (37) into (31), we obtain the reduced mean and fluctuating vertical component of the momentum equation at leading order,

$$\frac{\partial \bar{p}_{00}}{\partial \zeta} = 0, \quad (42a)$$

$$\frac{\partial w'_{01}}{\partial t_f} + \bar{\mathbf{u}}_{\perp 00} \cdot \nabla_f w'_{01} = -\frac{\partial p'_{01}}{\partial \zeta} + b'_{03} + \frac{\alpha}{Re_b} \left(\nabla_f^2 w'_{01} + \frac{\partial^2 w'_{01}}{\partial \zeta^2} \right), \quad (42b)$$

where the higher-order viscous term is added to regularize the fluctuation equation.

The set of equations formed by (38), (39), (41), and (42) are the intermediate regime analogs of the reduced model given in Chini et al. (2022). They are valid as long as $Re_b \geq O(1)$, and $\alpha \ll Pe_b \ll 1$. Given that $\alpha = Fr$ in this regime, this inequality constraint is equivalent to requiring that $Fr \ll 1$ (so $\alpha \ll 1$), $Re \geq Fr^{-2}$, and $Fr^{-1} \ll Pe \ll Fr^{-2}$. The physical implication of these equations, and their potential caveats, are discussed in § 4.

3.3.2 Case 2: $Pe_b \ll \alpha$ (the fully diffusive regime)

We now consider the regime where $Pe_b \ll \alpha$, in which both mean and fluctuating buoyancy fields are dominated by diffusion. Accordingly, we refer to this part of parameter space as the fully diffusive regime. Inspection of (30b) shows that the diffusion term in the fluctuation equation is unbalanced unless $b' = O(Pe_b w')$. We previously found that the vertical velocity fluctuations are $O(\alpha^{1/2})$, which implies here that $b' = O(\alpha^{1/2} Pe_b)$. We conclude that $b'_{0i} = 0 \forall i$, and that $b'_{10} = 0$ as well. Combined with the results obtained from analysis of the mean buoyancy equation in the diffusive limit, we conclude that $b_{10} = 0$ while $b_{11} = b'_{11}$.

Using this information and substituting (37) into (30), we obtain at lowest order

$$\bar{w}_{02} = \frac{\partial^2 \bar{b}_{12}}{\partial \zeta^2}, \quad (43a)$$

$$w'_{01} = \nabla_f^2 b'_{11} + \frac{\partial^2 b'_{11}}{\partial \zeta^2}, \quad (43b)$$

which is as expected from the LPN dynamics central to this regime (Lignières, 1999). The buoyancy equation is linear and does not contain any time dependence, instead instantaneously coupling the vertical velocity

and buoyancy fields to one another. The validity of Lignières' LPN equations was verified numerically by Cope et al. (2020), for instance.

As usual, the last step of the derivation involves analysis of the vertical component of the momentum equation. As in § 3.3.1, requiring hydrostatic equilibrium for the mean flow would imply $\alpha^2 = Fr^2/Pe_b$ (Lignières, 2020; Skoutnev, 2023), but leads to an inconsistency in the fluctuation equation, where the buoyancy term would be unbalanced. To see this, note that $b'/Fr^2 = O(\alpha^{1/2}Pe_b/Fr^2) = O(\alpha^{-3/2})$ with that choice for α , while the fluctuating pressure term (and all other dominant terms in the equation) is only $O(\alpha^{-1/2})$. As in the previous section, the resolution to this inconsistency is to insist that the buoyancy term in the fluctuation equation should be balanced instead. Here, this implies

$$O\left(\frac{p'}{\alpha}\right) = O\left(\frac{b'}{Fr^2}\right) \rightarrow \alpha = \frac{Fr^2}{Pe_b}, \quad (44)$$

using the fact that $p' = O(\alpha^{1/2})$ and $b' = O(\alpha^{1/2}Pe_b)$. Recalling that $Pe_b = \alpha^2Pe$, we then recover the crucial scaling relationship

$$\alpha = \left(\frac{Fr^2}{Pe}\right)^{1/3} = Fr_M^{4/3}, \quad (45)$$

which had originally been proposed by Cope et al. (2020) based on their DNS data. Our multiscale analysis therefore provides a sound theoretical basis for their empirical results.

After substituting (37) into (31) and using the available information, we obtain

$$\frac{\partial \bar{p}_{00}}{\partial \zeta} = 0, \quad (46a)$$

$$\frac{\partial w'_{01}}{\partial t_f} + (\bar{\mathbf{u}}_{\perp 00} \cdot \nabla_f)w'_{01} = -\frac{\partial p'_{01}}{\partial \zeta} + b'_{11} + \frac{\alpha}{Re_b} \left(\nabla_f^2 w'_{01} + \frac{\partial^2 w'_{01}}{\partial \zeta^2} \right), \quad (46b)$$

which is very similar to the system obtained in the intermediate regime studied in § 3.3.1, except for the appearance of the buoyancy fluctuation term b'_{11} instead of b'_{03} . As before, formally higher-order viscous terms are retained to regularize the fluctuation equation.

The set of equations formed by (38), (39), (43), and (46) are the fully-diffusive regime analogs of the reduced model derived by Chini et al. (2022). They are valid as long as $Re_b \geq O(1)$, and $Pe_b \ll \alpha \ll 1$. Given that $\alpha = (Fr^2/Pe)^{1/3}$ in this regime, this parameter constraint is equivalent to requiring that $Fr^2 \ll Pe$ (to ensure $\alpha \ll 1$), $Pe \ll Fr^{-1}$ (to ensure $Pe_b \ll \alpha$), and $Pe \geq Pr^3 Fr^{-4}$ (to ensure that $Re_b \geq O(1)$). The three conditions demarcate a triangle in logarithmic parameter space in which the equations are valid – see § 4.2. As an important self-consistency check, we see that the $Pe_b = \alpha$ transition between the fully diffusive and intermediate regimes is the same ($Pe = Fr^{-1}$) whether the transition is approached from the former or latter part of parameter space. In the next section (§ 4), we consider the physical implications of these equations as well as potential caveats on their applicability.

4 Discussion

In this study, we have extended the results of Chini et al. (2022) by performing a multiscale asymptotic analysis of stratified turbulence at low Prandtl number. Our work demonstrates the existence of several different regimes depending on the strength of the stratification (quantified by the inverse Froude number) and the rate of buoyancy diffusion (quantified by the inverse Péclet number). In each regime, the asymptotic analysis self-consistently yields a slow-fast system of quasilinear equations describing the concurrent evolution of a highly anisotropic, slow, large-scale mean flow and isotropic, fast, small-scale fluctuations. The large-scale anisotropy is characterized by the aspect ratio α (the ratio of the vertical to horizontal scales of the large-scale flow), whose functional dependence on Fr and Pe naturally emerges from the analysis. The various regime boundaries are locations in parameter space where relevant Reynolds and Péclet numbers arising in the mean flow and fluctuation equations are $O(1)$, signifying transitions between viscous and (formally) inviscid dynamics, and/or diffusive and non-diffusive dynamics. We now summarize our findings in each regime and then discuss the model assumptions as well as the implications of our results for low Prandtl number fluids.

4.1 Synopsis of multiscale equations and their validity

The first regime is characterized by $Fr \ll 1$ and $Pe_b, Re_b \geq O(1)$, where $Re_b = \alpha^2 Re$ and $Pe_b = \alpha^2 Pe$. In this regime, we recover the reduced model of Chini et al. (2022) and confirm that $\alpha = Fr$. Recalling that, at leading order, $w'_1 = w'/\alpha^{1/2}$, $b'_3 = b'/\alpha^{3/2}$, etc., that $\partial/\partial t_f = \alpha\partial/\partial t$ (and similarly for ∇_f and ∇_\perp), and finally that $\partial/\partial \zeta = \alpha\partial/\partial z$, we can rewrite (33), (34), (35) and (36) as the following quasilinear system for mean and fluctuations, expressed in the original *isotropic* (single-scale) variables.

Mean flow equations:

$$\frac{\partial \bar{\mathbf{u}}_\perp}{\partial t} + \bar{\mathbf{u}}_\perp \cdot \nabla \bar{\mathbf{u}}_\perp = -\nabla_\perp \bar{p} - \frac{\partial}{\partial z} (\overline{w' \mathbf{u}'_\perp}) + \frac{1}{Re} \frac{\partial^2 \bar{\mathbf{u}}_\perp}{\partial z^2} + \bar{F} \hat{\mathbf{e}}_x, \quad (47a)$$

$$\frac{\partial \bar{p}}{\partial z} = \frac{\bar{b}}{Fr^2}, \quad \nabla \cdot \bar{\mathbf{u}} = 0, \quad (47b)$$

$$\frac{\partial \bar{b}}{\partial t} + \bar{\mathbf{u}}_\perp \cdot \nabla \bar{b} + \bar{w} = -\frac{\partial}{\partial z} (\overline{w' b'}) + \frac{1}{Pe} \frac{\partial^2 \bar{b}}{\partial z^2}. \quad (47c)$$

Fluctuation equations:

$$\frac{\partial \mathbf{u}'_\perp}{\partial t} + \bar{\mathbf{u}}_\perp \cdot \nabla_\perp \mathbf{u}'_\perp + w' \frac{\partial \bar{\mathbf{u}}_\perp}{\partial z} = -\nabla_\perp p' + \frac{1}{Re} \nabla^2 \mathbf{u}'_\perp, \quad (47d)$$

$$\frac{\partial w'}{\partial t} + \bar{\mathbf{u}}_\perp \cdot \nabla_\perp w' = -\frac{\partial p'}{\partial z} + \frac{b'}{Fr^2} + \frac{1}{Re} \nabla^2 w', \quad (47e)$$

$$\nabla \cdot \mathbf{u}' = 0, \quad (47f)$$

$$\frac{\partial b'}{\partial t} + \bar{\mathbf{u}}_\perp \cdot \nabla_\perp b' + w' \frac{\partial \bar{b}}{\partial z} + w' = \frac{1}{Pe} \nabla^2 b'. \quad (47g)$$

This system, which is equivalent to equations (2.28)–(2.35) in Chini et al. (2022), is valid when $Fr \ll 1$, and $Re, Pe \geq O(Fr^{-2})$, corresponding to $Re_b, Pe_b \geq O(1)$. One advantage of re-casting the equations in the original isotropic variables is now clear: the system has the exact form that one would, absent the asymptotic analysis, *assume* when making a quasilinear reduction of the Boussinesq equations (e.g. Garaud, 2001; Fitzgerald and Farrell, 2018, 2019). In this work, we have demonstrated that this form is in fact a natural outcome of the slow-fast asymptotic expansion, and is therefore asymptotically exact in the given distinguished limit.

Dimensionally, this regime has a characteristic vertical scale $H^* = \alpha L^* = U^*/N^*$. The characteristic vertical velocity is $W^* = \alpha^{1/2} U^* = (U^{*3}/N^* L^*)^{1/2}$, and w is dominated by small-scale fluctuations. The characteristic buoyancy scale is $N^{*2} H^*$, and b and dominated by large scales. Note that the horizontal field u is predominantly large scale, by assumption, but contains small scales with (dimensional) amplitudes $O(Fr^{1/2} U^*)$. These findings have been validated in numerical and laboratory experiments (Holford and Linden, 1999; Brethouwer et al., 2007; Oglethorpe et al., 2013; Maffioli and Davidson, 2016), with at least suggestive observational evidence of this ‘layered anisotropic stratified turbulence’ (LAST) regime being obtainable from seismic oceanography surveys (Falder et al., 2016).

The scenario in which $Fr \ll 1$, $Re_b \geq O(1)$ and $\alpha \ll Pe_b \ll 1$ is an intermediate regime where the mean flow is diffusive while the fluctuations are not. In this regime, we also find that $\alpha = Fr$. The slow-fast system of equations (38), (39), (41), and (42), expressed in the original isotropic single-scale variables, becomes the following quasilinear system.

Mean flow equations:

$$\frac{\partial \bar{\mathbf{u}}_\perp}{\partial t} + \bar{\mathbf{u}}_\perp \cdot \nabla \bar{\mathbf{u}}_\perp = -\nabla_\perp \bar{p} - \frac{\partial}{\partial z} (\overline{w' \mathbf{u}'_\perp}) + \frac{1}{Re} \frac{\partial^2 \bar{\mathbf{u}}_\perp}{\partial z^2} + \bar{F} \hat{\mathbf{e}}_x, \quad (48a)$$

$$\frac{\partial \bar{p}}{\partial z} = 0, \quad \nabla \cdot \bar{\mathbf{u}} = 0, \quad (48b,c)$$

$$\bar{w} = -\frac{\partial}{\partial z} (\overline{w' b'}) + \frac{1}{Pe} \frac{\partial^2 \bar{b}}{\partial z^2}. \quad (48d)$$

Fluctuation equations:

$$\frac{\partial \mathbf{u}'_\perp}{\partial t} + \bar{\mathbf{u}}_\perp \cdot \nabla_\perp \mathbf{u}'_\perp + w' \frac{\partial \bar{\mathbf{u}}_\perp}{\partial z} = -\nabla_\perp p' + \frac{1}{Re} \nabla^2 \mathbf{u}'_\perp, \quad (48b)$$

$$\frac{\partial w'}{\partial t} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_{\perp} w' = -\frac{\partial p'}{\partial z} + \frac{b'}{Fr^2} + \frac{1}{Re} \nabla^2 w', \quad (48c)$$

$$\nabla \cdot \mathbf{u}' = 0, \quad (48d)$$

$$\frac{\partial b'}{\partial t} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_{\perp} b' + w' = \frac{1}{Pe} \nabla^2 b'. \quad (48e)$$

These equations are valid for $Re \geq Fr^{-2}$ (so $Re_b \geq O(1)$) and $Fr^{-1} \ll Pe \ll Fr^{-2}$ (so $\alpha \ll Pe_b \ll 1$). By writing them in the original isotropic variables, we now see that the correct quasilinear equations at this order are almost the same as in the Chini et al. (2022) regime, except that terms in \bar{b} are dropped (in the mean hydrostatic balance and in the buoyancy perturbation equation) because they are formally of higher order. In addition, the mean buoyancy equation takes the LPN form of Lignières et al. (1999), modified by the fluctuation-induced buoyancy flux.

Dimensionally, the vertical length scale and vertical velocity scale are the same as in the non-diffusive regime. The buoyancy field, however, is now dominated by large scales if $Pe_b \geq \alpha^{1/2}$ and by small scales if $Pe_b \leq \alpha^{1/2}$. Testing these scaling laws numerically will be very difficult, unfortunately, because the range of the intermediate region is very small (holding Pe constant while varying Fr^{-1} or vice-versa) unless Pr is itself very small.

Finally, the case where $Fr \ll 1$, $Re_b \geq O(1)$ and $Pe_b \ll \alpha$ corresponds to a fully diffusive regime in which both the mean flow and fluctuations are dominated by diffusion and satisfy the LPN balance derived by Lignières (1999). In this regime, we have demonstrated that $\alpha = (Fr^2/Pe)^{1/3}$. The slow-fast system of equations (38), (39), (43) and (46), written in the original isotropic single-scale variables, is given below.

Mean flow equations:

$$\frac{\partial \bar{\mathbf{u}}_{\perp}}{\partial t} + \bar{\mathbf{u}}_{\perp} \cdot \nabla \bar{\mathbf{u}}_{\perp} = -\nabla_{\perp} \bar{p} - \frac{\partial}{\partial z} (\overline{w' \mathbf{u}'_{\perp}}) + \frac{1}{Re} \frac{\partial^2 \bar{\mathbf{u}}_{\perp}}{\partial z^2} + \bar{F} \hat{\mathbf{e}}_x, \quad (49a)$$

$$\frac{\partial \bar{p}}{\partial z} = 0, \quad \nabla \cdot \bar{\mathbf{u}} = 0, \quad (49b,c)$$

$$\bar{w} = \frac{1}{Pe} \frac{\partial^2 \bar{b}}{\partial z^2}. \quad (49d)$$

Fluctuation equations:

$$\frac{\partial \mathbf{u}'_{\perp}}{\partial t} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_{\perp} \mathbf{u}'_{\perp} + w' \frac{\partial \bar{\mathbf{u}}_{\perp}}{\partial z} = -\nabla_{\perp} p' + \frac{1}{Re} \nabla^2 \mathbf{u}'_{\perp}, \quad (49b)$$

$$\frac{\partial w'}{\partial t} + \bar{\mathbf{u}}_{\perp} \cdot \nabla_{\perp} w' = -\frac{\partial p'}{\partial z} + \frac{b'}{Fr^2} + \frac{1}{Re} \nabla^2 w', \quad (49c)$$

$$\nabla \cdot \mathbf{u}' = 0, \quad (49d)$$

$$w' = \frac{1}{Pe} \nabla^2 b'. \quad (49e)$$

This system is valid for $Re_b \geq O(1)$, and $Pe_b \ll \alpha \ll 1$. As $\alpha = (Fr^2/Pe)^{1/3}$, these constraints are equivalent to $Fr^2 \ll Pe$ (so that $\alpha \ll 1$), $Pe \ll Fr^{-1}$ (so that $Pe_b \ll \alpha$), and $Pe \geq Pr^3 Fr^{-4}$ (so that $Re_b \geq O(1)$).

Dimensionally, the characteristic vertical length scale $H^* = \alpha L^* = (Fr^2/Pe)^{1/3} L^* = (U^* \kappa^*/N^{*2})^{1/3}$. The characteristic vertical velocity $W^* = \alpha^{1/2} U^* = (Fr^2/Pe)^{1/6} U^* = (U^{*7} \kappa^*/N^{*2} L^{*3})^{1/6}$, and is dominated by small-scale fluctuations. The characteristic buoyancy scale is $Pe(Fr^2/Pe)^{5/6} L^* N^{*2} = (U^{*11/6} \kappa^{*-1/6} N^{*-5/3} L^{*-3/2}) L^* N^{*2}$ and is dominated by small-scale fluctuations as well. These scalings have been validated by the DNS of Cope et al. (2020).

Crucially, we find that in all three regimes the characteristic vertical velocity $W^* = \alpha^{1/2} U^*$ is significantly larger than that predicted from the anisotropically scaled equations given in § 2.1 and § 2.2, where $W^* = \alpha U^*$. This larger scaling has implications for turbulent vertical transport of buoyancy and passive scalars in stellar interiors (see § 5).

Finally, note that in all of these regimes, we have assumed $Re_b \geq O(1)$ which then implies that $Re_b \gg \alpha$ since $\alpha \ll 1$. Recalling that the fluctuation equations have an effective Reynolds number Re_b/α , this condition is necessary to ensure that the small-scale fluctuations can develop without being suppressed by

viscosity, and is therefore key to the multiscale expansions derived here. When $Re_b < 1$, viscous effects become important and could strongly affect our conclusions. We do not pursue this issue further here, deferring discussion of viscous regimes to a future publication.

However, our results thus shed doubt on whether there is any region of parameter space in which the anisotropically-scaled equations (see § 2.1 in the large Pr limit and § 2.2 in the low Pr limit) apply while viscous effects are at the same time negligible. Indeed, by construction these equations are valid only when small-scale fluctuations are suppressed, but we see that this requires $Re_b \leq O(\alpha)$, in which case viscosity should play a dominant role in the mean equations and would impart a vertical length scale to the mean flow that necessarily is $O(Re^{-1/2})$ (c.f. Godoy-Diana et al., 2004; Brethouwer et al., 2007; Waite, 2014; Cope et al., 2020).

Using existing algorithms for slow-fast systems (Michel and Chini, 2019; Chini et al., 2022; Ferraro, 2022), our ongoing work involves numerically solving these multiscale equations for a range of values of Re_b , which would allow numerical identification of values at which these behaviours begin to take effect (Shah, 2022).

4.2 Regimes of stratified stellar turbulence

Our findings partition parameter space into various regimes of stratified turbulence, which are illustrated in figure 1 for a $Pr = 1$ fluid (top) and a $Pr = 10^{-6}$ fluid (bottom). The latter is typical of some stellar interiors (see Garaud, 2021). In both panels, the horizontal axis shows the inverse Froude number Fr^{-1} , so that stratification increases to the right. The vertical axis shows the outer scale Péclet number Pe , so that diffusive effects decrease upward. In both panels, the grey region shows where α is $O(1)$, so the large-scale flow is isotropic; this regime is not the focus of our study. For stronger stratification, the large-scale flow becomes anisotropic, and the possible regimes of stratified turbulence depend on Pr .

For $Pr = O(1)$, shown in the upper plot, the partitioning of parameter space for $Fr^{-1} \gg 1$ is straightforward. When $Pe_b, Re_b \geq O(1)$ (green region), viscosity and diffusion play a secondary role in both the mean and fluctuation dynamics. This is the regime studied by Chini et al. (2022). By contrast, if $Pe_b, Re_b \ll 1$ (white region) then both effects begin to influence the flow dynamics and cannot be ignored. The transition takes place when $Re, Pe \approx Fr^{-2}$.

At low Pr , diffusion becomes important long before viscosity does, so the $Pe_b = O(1)$ transition is distinct from the $Re_b = O(1)$ transition. This distinction opens up parameter space to the two new regimes discussed in this work: the intermediate regime (yellow region, the dynamics of which are described in § 3.3.1) and the fully-diffusive regime (purple region, the dynamics of which are described in § 3.3.2). We now clearly see that the fully-diffusive regime is confined to a triangle in log-log parameter space for a given Prandtl number. It is delimited from above by the intermediate regime (yellow region), from below by the isotropic regime (grey region), and from the right by the viscous regime (white region). More specifically, it is bounded by the points A ($Fr^{-1} = 1, Pe = 1$), B ($Fr^{-1} = Pr^{-1}, Pe = Pr^{-1}$) and C ($Fr^{-1} = Pr^{-1/2}, Pe = Pr$) and thus becomes increasingly wide as Pr decreases, but shrinks towards the point A as $Pr \rightarrow 1$.

A major implication of our results is that, depending on the choice of Pe , different regimes are encountered as the stratification increases. We now discuss horizontal transects through figure 1, for different values of Pe . We first consider the case $Pr < Pe < 1$, which is the regime discussed in Cope et al. (2020). As Fr^{-1} increases, our model predicts that the turbulence ought to be isotropic until $Fr^{-1} = Pe^{-1/2}$ (interestingly, because diffusion partially relaxes the effects of stratification when $Pe \ll 1$). As Fr^{-1} continues to increase, the turbulence enters the fully diffusive (anisotropic) regime, and remains in that regime until $Fr^{-1} = (Pe/Pr^3)^{1/4}$, at which point viscosity begins to affect the mean flow. This series of regime transitions is qualitatively consistent with what is observed in the low Prandtl number DNS of Cope et al. (2020).

At the other extreme, let us consider a transect for $Pe > Pr^{-1}$, which is the case considered by Garaud (2020), who primarily analysed simulations for which $Pe = 60$ and $Pr = 0.1$. This regime is relevant for strongly sheared layers in stellar interiors, such as the solar tachocline. For moderate stratification, namely $1 \leq Fr^{-1} \leq Pe$, our analysis shows that the turbulence is expected to be both anisotropic and non-diffusive, and its properties should be captured by the model of Chini et al. (2022). As stratification increases past $Fr^{-1} = Pe$, the turbulence is predicted to enter the intermediate regime, where the mean flow is dominated by diffusion but the fluctuations are not. Beyond $Fr^{-1} = \sqrt{Pe/Pr}$, viscous effects should become important. Note how, at these large values of the Péclet number, the fully diffusive inviscid regime discussed in § 3.3.2 is not accessible. Instead, viscosity begins to influence the mean flow before diffusion




Figure 1: Regime diagram for a fluid with $Pr = 1$ (top) and a fluid with $Pr = 10^{-6}$ (bottom). The Péclet number is on the vertical axis and the inverse Froude number is on the horizontal axis (such that stratification increases to the right). Regions of unstratified, isotropic turbulence are marked in grey. Regions in white are viscously controlled, and will be discussed in future work. In both panels, the blue dashed line marks the transition $Pe_b = O(1)$ where the mean flow becomes diffusive, and the red solid lines mark the viscous transition where $Re_b = O(1)$. With $Pr \ll 1$ (bottom panel), it is possible to have $Pe_b \ll 1 \ll Re_b$, i.e. regimes of thermally diffusive stratified turbulence, which are not possible when $Pr = O(1)$. In that case, the solid blue line marks the transition where $Pe_b = O(\alpha)$ where the small-scale fluctuations become diffusive. A region of parameter space opens up between $Pe_b = O(1)$ and $Re_b = O(1)$, where the two new regimes identified in this work exist: the intermediate regime, marked in yellow, and the fully diffusive regime marked in purple, bounded by the three corners of the triangle labelled A, B, C (see main text). A parameter set indicative of the solar tachocline is marked with a black circle, using typical parameters given by Garaud (2020) ($Pe = 10^8$, $Fr^{-1} = 10^3$).

influences the fluctuations.

This series of regime transitions is indeed qualitatively consistent with the simulations reported in Garaud (2020). Specifically, she found that the turbulence is mostly isotropic for $Fr^{-1} < 1$, then becomes anisotropic with little effect of diffusion for intermediate values of Fr^{-1} . However, her empirically-derived scaling laws ($H^* \propto Fr^{2/3}L^*$, $W^* \propto Fr^{2/3}U^*$) do not match those predicted by the Chini et al. (2022) theory; a target for future work is to revisit the data to determine whether this discrepancy can be explained. For even larger values of Fr^{-1} , Garaud (2020) found that diffusion becomes important, and that the dynamics are governed by the LPN equations. However, at this point the turbulence is also in an intermittent regime where viscosity partially suppresses the fluctuations, a situation qualitatively consistent with predictions from figure 1.

Finally, the case where $1 < Pe < Pr^{-1}$ is relevant for weaker shear layers in stellar interiors. As Fr^{-1} increases, our multiscale analysis predicts that the turbulence ought to be isotropic until $Fr^{-1} = 1$, at which point the regime analysed by Chini et al. (2022) sets in and both the mean and fluctuation dynamics are non-diffusive. As the stratification continues to increase, the vertical eddy scale decreases gradually until the mean flow becomes dominated by diffusion at $Fr^{-1} = \sqrt{Pe}$ and the intermediate regime is manifest. The flow dynamics remain essentially unchanged in that regime until diffusion also begins to affect the fluctuations as well, at $Fr^{-1} = Pe$, at which point the turbulence becomes fully diffusive. Finally, viscous effects begin to be important when $Fr^{-1} = (Pe/Pr^3)^{1/4}$. To date, no DNS have been published probing this range of Pe . Verifying the existence of these successive transitions with DNS is one of our areas of active work.

4.3 Mathematical considerations

Having constructed a map of parameter space based on the results of our multiscale analysis, we now discuss important consequences and caveats of the resulting reduced models. As in Chini et al. (2022), we find that the reduced equations in each regime form a closed set describing the concurrent evolution of a highly anisotropic large-scale mean flow together with isotropic small-scale fluctuations. Crucially, the fluctuation equations derived in all regimes identified are linear in the fluctuation fields, but feed back nonlinearly on the mean flow evolution. The reduced models thus all have a quasilinear form, which emerges self-consistently from the asymptotic analysis rather than being imposed *a priori*. This emergent quasilinearity in the slow-fast limit therefore appears to be an inherent property of stratified turbulence regardless of the Prandtl number. Our findings can be used as a theoretical basis not only for using the quasilinear approximation to create reduced models of stratified turbulence, but also to know precisely which terms to keep in each region of parameter space (see, e.g, Marston and Tobias, 2023). Of course, the fundamental premise upon which the analysis is predicated is that the stratification drives scale-separated dynamics characterized by the anisotropic large-scale and roughly isotropic small-scale flow structures. To the extent that spectrally non-local interactions between these disparate scales of motion play a crucial role in stratified turbulence, the conclusions and predictions of our multiscale analysis are likely to be valid.

In quasilinear systems subject to fast instabilities, nonlinear saturation requires the feedback from the finite-amplitude fluctuations to render the mean fields marginally stable to disturbances of any horizontal wavenumber. It is straightforward to verify that this condition of approximate marginal stability indeed characterizes each regime. This observation is consistent with the empirical observation of ‘self-organized criticality’ mentioned in §1, giving further credence to the appropriateness of the multi-scale approach used here. In the $Pe_b \geq O(1)$ and intermediate regimes, the fluctuation equations are non-diffusive at leading order, and therefore describe the growth (or decay) of perturbations due to a standard vertical shear instability. The dimensional vertical shear S^* of the mean flow can be estimated to be roughly U^*/H^* since $|\bar{\mathbf{u}}| = O(1)$. The gradient Richardson number

$$J = \frac{N^{*2}}{S^{*2}} = O\left(\frac{N^{*2}H^{*2}}{U^{*2}}\right) = O\left(\frac{\alpha^2}{Fr^2}\right) = O(1) \quad (50)$$

in these regimes, which is indeed marginally-stable to the Richardson and Miles–Howard criteria (Richardson, 1920; Miles, 1961; Howard, 1961). Note that Garaud et al. (2023) recently verified that J is indeed $O(1)$ in DNS of stratified turbulence at $Pe_b = O(1)$.

In the $Pe_b \ll \alpha$ regime, by contrast, the fluctuation equations are inherently diffusive, and one therefore expects the vertical shear instability to be of the diffusive kind (Zahn, 1974; Jones, 1977; Lignières et al.,

1999). It has been shown, at least for sinusoidal shear flows (Garaud et al., 2015a), that the condition for marginal linear stability is $JPe_S = O(1)$ where $Pe_S = U^*H^*/\kappa^*$ is the Péclet number based on the vertical shear profile. We can easily check that the scalings found in § 3.3.2 satisfy this criterion. Indeed,

$$JPe_S = O\left(\frac{N^{*2}H^{*2}U^*H^*}{U^{*2}\kappa^*}\right) = O\left(\alpha^3\frac{Pe}{Fr^2}\right) = O(1), \quad (51)$$

as required. Taken together with the empirical scaling laws obtained from the DNS data of Cope et al. (2020), this evidence confirms that (45) is indeed the correct expression for α in stratified turbulence at very low Pe_b .

A somewhat less intuitive result of our analysis is the requirement that $\partial\bar{p}_{00}/\partial\zeta = 0$, while $\bar{p}_{00} = O(1)$, in both intermediate and fully diffusive regimes. Physically, this condition can be understood by noting that in the $Pe_b \ll 1$ scenario, departures from the mean background stratification formally are exceedingly small (owing to the strong thermal diffusion), which explains why they do not affect the assumed *background* hydrostatic balance. Furthermore, recalling that ζ has been rescaled by the vertical length scale $H^* = \alpha L^*$ with $\alpha \ll 1$, we note that $\partial_\zeta\bar{p}_{00} = 0$ only applies on that scale, and does not preclude \bar{p}_{00} from potentially varying on larger scales. Indeed, the possibility exists that *two* vertical scales may be incorporated into the dynamics, with the leading-order mean pressure being hydrostatically coupled to the leading-order mean buoyancy on the larger vertical scale, although we leave exploration of that possibility for future work. Regardless, we emphasize that buoyancy anomalies *do* affect the fluctuation dynamics, which in turn modifies the mean flow.

5 Conclusions

In this study, inspired by numerical evidence of scale separation and flow anisotropy, we have conducted a formal multiscale asymptotic analysis of the Boussinesq equations governing the dynamics of strongly stratified turbulence at low Prandtl number. A key outcome of our work is a new map of parameter space, shown in figure 1, that demarcates different regimes of stratified turbulence. Crucially, we find that new regions of parameter space open up at low Prandtl number in which diffusive turbulent flows exist, scenarios that are not possible at $Pr = O(1)$. For each of these new regimes, scaling laws for the vertical velocity and vertical length scale of turbulent eddies naturally emerge from the analysis. These scaling laws are summarized in § 4.2 and recover previous findings by Chini et al. (2022) and Cope et al. (2020) in appropriate distinguished limits. Finally, recent work by Garaud et al. (2023) has demonstrated numerically that the presence of a mean vertical shear (ignored in this work) has little impact on these scalings as long as its amplitude is smaller than the small-scale emergent vertical shear U^*/H^* . This finding can be proved more formally using the asymptotic tools developed here, suggesting that the new regime diagram is robust (at least if other effects such as rotation and magnetic fields are absent; see below).

These results have important implications for low Pr fluids, such as liquid metals (where $Pr \sim 0.01 - 0.1$), planetary interiors (where $Pr \sim 0.001 - 0.1$) and stellar interiors (where $Pr \sim 10^{-9} - 10^{-2}$). In particular, the scaling laws derived enable us to propose simple parameterizations for the turbulent diffusion coefficient (D^*) in each identified regime by multiplying the characteristic vertical length and vertical velocity scales.

- In the $Pe_b \geq O(1)$ and the intermediate regimes,

$$D^* \propto H^*W^* \propto Fr^{3/2}L^*U^* \propto U^{*5/2}/(L^{*1/2}N^{*3/2}). \quad (52)$$

- In the fully diffusive regime,

$$D^* \propto H^*W^* \propto (Fr^2/Pe)^{1/2}L^*U^* \propto U^{*3/2}\kappa^{*1/2}/(L^{*1/2}N^*). \quad (53)$$

As such, our work challenges current understanding of stratified turbulence in stars. Indeed, the most commonly used model for stratified turbulence in stellar evolution calculations is the model of Zahn (1992), which proposes a vertical turbulent diffusivity equivalent to (53). However, we have demonstrated that this expression is only valid in a relatively small region of the parameter space (see figure 1). In particular, Zahn’s model assumes that the turbulence is always fully diffusive, but this is not the case in the intermediate and

$Pe_b \geq O(1)$ regimes, which are appropriate for more strongly sheared fluid layers such as the solar tachocline. Zahn’s model also assumes that viscosity is negligible, which is not the case for sufficiently stratified flows.

Of course, future work will be needed to investigate the effects of rotation and magnetization, which are also important in stellar and planetary interiors. Both effects are likely to stabilize the horizontal turbulence to some extent, which will therefore affect the emergent vertical shear instability as well. Additionally, we need to systematically compare the model predictions with existing DNS (Brethouwer et al., 2007; Maffioli and Davidson, 2016; Cope et al., 2020; Garaud, 2020) as well as with future DNS conducted at more extreme values of the Reynolds, Prandtl, and Péclet numbers. In particular, known discrepancies between the model and data need to be explained (see § 4.2). This step is crucial to gain enough confidence in the model to apply it at stellar and planetary parameter values. Nonetheless, this work has already demonstrated the power of formal multiscale asymptotic analysis for discovering and validating the existence of new regimes of stratified turbulence in stellar and planetary interiors.

Acknowledgements

The authors gratefully acknowledge the Geophysical Fluid Dynamics Summer School (NSF 1829864), particularly the 2022 program. P.G. acknowledges funding from NSF AST-1814327. K.S. acknowledges funding from the James S. McDonnell Foundation. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Declaration of Interests

The authors report no conflict of interest.

References

- Aerts, C., Mathis, S., and Rogers, T. (2019). Angular Momentum Transport in Stellar Interiors. *Ann. Rev. Ast. Astrophys.*, 57:35–78.
- Augier, P., Billant, P., and Chomaz, J.-M. (2015). Stratified turbulence forced with columnar dipoles: numerical study. *Journal of Fluid Mechanics*, 769:403–443.
- Bartello, P. and Tobias, S. (2013). Sensitivity of stratified turbulence to the buoyancy Reynolds number. *Journal of Fluid Mechanics*, 725:1–22.
- Basak, S. and Sarkar, S. (2006). Dynamics of a stratified shear layer with horizontal shear. *J. Fluid Mech.*, 568:19–54.
- Batchelor, G. (1967). *An introduction to fluid dynamics*. Cambridge university press.
- Billant, P. and Chomaz, J.-M. (2000). Experimental evidence for a new instability of a vertical columnar vortex pair in a strongly stratified fluid. *Journal of Fluid Mechanics*, 418(1):167–188.
- Billant, P. and Chomaz, J.-M. (2001). Self-similarity of strongly stratified inviscid flows. *Physics of Fluids*, 13(6):1645–1651.
- Brethouwer, G., Billant, P., Lindborg, E., and Chomaz, J.-M. (2007). Scaling analysis and simulation of strongly stratified turbulent flows. *Journal of Fluid Mechanics*, 585:343–368.
- Caulfield, C. (2021). Layering, instabilities and mixing in turbulent stratified flow. *Annu. Rev. Fluid Mech.*, 53:113–145.
- Chini, G., Michel, G., Julien, K., Rocha, C., and Caulfield, C. (2022). Exploiting self-organized criticality in strongly stratified turbulence. *Journal of Fluid Mechanics*, 933.

- Cope, L., Garaud, P., and Caulfield, C. (2020). The dynamics of stratified horizontal shear flows at low Péclet number. *J. Fluid Mech.*, 903:A1.
- Dudis, J. (1973). The stability of a thermally radiating stratified shear layer. *Journal of Fluid Mechanics*, 58:81–96.
- Falder, M., White, N. J., and Caulfield, C. P. (2016). Seismic imaging of rapid onset of stratified turbulence in the South Atlantic Ocean. *J. Phys. Oceanogr.*, 46(4):1023–1044.
- Ferraro, A. (2022). Exploiting marginal stability in slow-fast quasilinear dynamical systems. Technical report, EPFL.
- Fitzgerald, J. and Farrell, B. (2018). Statistical state dynamics of vertically sheared horizontal flows in two-dimensional stratified turbulence. *Journal of Fluid Mechanics*, 854:544–590.
- Fitzgerald, J. and Farrell, B. (2019). Statistical state dynamics analysis of buoyancy layer formation via the phillips mechanism in two-dimensional stratified turbulence. *Journal of Fluid Mechanics*, 864:R3.
- Garaud, P. (2001). Latitudinal shear instability in the solar tachocline. *MNRAS*, 324(1):68–76.
- Garaud, P. (2020). Horizontal shear instabilities at low Prandtl number. *The Astrophysical Journal*, 901(2).
- Garaud, P. (2021). Journey to the center of stars: The realm of low prandtl number fluid dynamics. *Physical Review Fluids*, 6(3):030501.
- Garaud, P., Gallet, B., and Bischoff, T. (2015a). The stability of stratified spatially periodic shear flows at low Péclet number. *Physics of Fluids*, 27(8):084104 1–21.
- Garaud, P., Khan, S., and Brown, J. (2023). The combined effects of vertical and horizontal shear instabilities. *arXiv e-prints*, page arXiv:2308.14442.
- Garaud, P., Medrano, M., Brown, J., Mankovich, C., and Moore, K. (2015b). Excitation of Gravity Waves by Fingering Convection, and the Formation of Compositional Staircases in Stellar Interiors. *ApJ*, 808(1):89.
- Godoy-Diana, R., Chomaz, J.-M., and Billant, P. (2004). Vertical length scale selection for pancake vortices in strongly stratified viscous fluids. *Journal of Fluid Mechanics*, 504:229–238.
- Gregg, M., D’Asaro, E., Riley, J., and Kunze, E. (2018). Mixing efficiency in the ocean. *Annu. Rev. Mar. Sci.*, 10(1):443–473.
- Holford, J. and Linden, P. (1999). Turbulent mixing in a stratified fluid. *Dynamics of Atmospheres and Oceans*, 30(2):173–198.
- Howard, L. (1961). Note on a paper of John W. Miles. *J. Fluid Mech.*, 10:509–512.
- Ivey, G., Winters, K., and Koseff, J. (2008). Density Stratification, Turbulence, but How Much Mixing? *Annual Review of Fluid Mechanics*, 40(1):169–184.
- Jackson, P. R. and Rehmann, C. R. (2014). Experiments on Differential Scalar Mixing in Turbulence in a Sheared, Stratified Flow. *J. Phys. Ocean.*, 44(10):2661–2680.
- Jacobitz, F. and Sarkar, S. (2000). A direct numerical study of transport and anisotropy in a stably stratified turbulent flow with uniform horizontal shear. *Flow, Turbul. Combust.*, 63:343–360.
- Jones, C. (1977). The Onset of Shear Instability in Stars. *Geophysical and Astrophysical Fluid Dynamics*, 8:165–184.
- Lignières, F. (1999). The small-Péclet-number approximation in stellar radiative zones. *Astronomy & Astrophysics*, 348:933–939.
- Lignières, F. (2020). Turbulence in stably stratified radiative zone. In *Multi-Dimensional Processes In Stellar Physics*, pages 111–140.

- Lignières, F., Califano, F., and Mangeney, A. (1999). Shear layer instability in a highly diffusive stably stratified atmosphere. *Astron. Astrophys.*, 349:1027–1036.
- Lucas, D., Caulfield, C., and Kerswell, R. (2017). Layer formation in horizontally forced stratified turbulence: connecting exact coherent structures to linear instabilities. *Journal of Fluid Mechanics*, 832:409–437.
- Maffioli, A. and Davidson, P. (2016). Dynamics of stratified turbulence decaying from a high buoyancy reynolds number. *Journal of Fluid Mechanics*, 786:210–233.
- Marston, J. and Tobias, S. (2023). Recent Developments in Theories of Inhomogeneous and Anisotropic Turbulence. *Annual Review of Fluid Mechanics*, 55:351–375.
- Mashayek, A., Caulfield, C., and Alford, M. H. (2022). Goldilocks mixing in oceanic shear-induced turbulent overturns. *J. Fluid Mech.*, 928:A1.
- Michel, G. and Chini, G. (2019). Multiple scales analysis of slow–fast quasi-linear systems. *Proceedings of the Royal Society A*, 475(2223):20180630.
- Miles, J. W. (1961). On the stability of heterogeneous shear flows. *Journal of Fluid Mechanics*, 10:496–508.
- Munk, W. (1966). Abyssal recipes. *Deep Sea Research A*, 13(4):707–730.
- Ogletorpe, R., Caulfield, C., and Woods, A. (2013). Spontaneous layering in stratified turbulent Taylor-Couette flow. *Journal of Fluid Mechanics*, 721:R3 1–12.
- Park, Y.-G., Whitehead, J., and Gnanadeskian, A. (1994). Turbulent mixing in stratified fluids: layer formation and energetics. *Journal of Fluid Mechanics*, 279:279–311.
- Pinsonneault, M. (1997). Mixing in Stars. *Ann. Rev. Ast. Astrophys.*, 35:557–605.
- Richardson, L. (1920). The Supply of Energy from and to Atmospheric Eddies. *Royal Society of London Proceedings Series A*, 97:354–373.
- Ruddick, B., McDougall, T., and Turner, J. (1989). The formation of layers in a uniformly stirred density gradient. *Deep Sea Research A*, 36(4):597–609.
- Salehipour, H., Peltier, W. R., and Caulfield, C. (2018). Self-organised criticality of turbulence in strongly stratified mixing layers. *J. Fluid Mech.*, 856:228–256.
- Shah, K. (2022). Scaling with the stars: the emergence of marginal stability in low pr turbulence. Technical report, Geophysical Fluid Dynamics Summer School 2022: Project Report.
- Skoutnev, V. (2023). Critical balance and scaling of strongly stratified turbulence at low Prandtl number. *Journal of Fluid Mechanics*, 956:A7.
- Smyth, W. D. and Moum, J. N. (2013). Marginal instability and deep cycle turbulence in the eastern equatorial pacific ocean. *Geophys. Res. Lett.*, 40(23):6181–6185.
- Smyth, W. D., Nash, J. D., and Moum, J. N. (2019). Self-organized criticality in geophysical turbulence. *Sci. Reports*, 9:Art. No. 3747.
- Spiegel, E. and Veronis, G. (1960). On the boussinesq approximation for a compressible fluid. *The Astrophysical Journal*, 131:442.
- Spiegel, E. and Zahn, J.-P. (1970). Instabilities of Differential Rotation. *Comments on Astrophysics and Space Physics*, 2:178.
- Townsend, A. (1958). The effects of radiative transfer on turbulent flow of a stratified fluid. *Journal of Fluid Mechanics*, 4:361–375.

- Waite, M. (2014). Direct numerical simulations of laboratory-scale stratified turbulence. *Modeling Atmospheric and Oceanic Flows: Insights from Laboratory Experiments and Numerical Simulations*, pages 159–175.
- Yi, Y. R. and Koseff, J. R. (2023). Underlying physics of mixing efficiency for shear-forced, stratified turbulence. *Phys. Rev. Fluids*, 8(8):084803.
- Zahn, J.-P. (1974). Rotational instabilities and stellar evolution. In *Symposium-International Astronomical Union*, volume 59, pages 185–195. Cambridge University Press.
- Zahn, J.-P. (1992). Circulation and turbulence in rotating stars. *Astronomy and Astrophysics*, 265:115–132.
- Zhou, Q. and Diamessis, P. (2019). Large-scale characteristics of stratified wake turbulence at varying reynolds number. *Phys. Rev. Fluids*, 4:084802.